



ISSN 2395-650X

**International Journal** of  
Life Sciences Biotechnology Pharma Sciences

IJLBPS



[www.ijlbps.org](http://www.ijlbps.org)

E-mail: [editorijlbps@gmail.com](mailto:editorijlbps@gmail.com) [editor@ijlbps.org](mailto:editor@ijlbps.org)

## Biosystematics studies on medicinal plant *Urginea indica* Kunth. liliaceae - A review

**Sethi Pooja**

### **Abstract**

*Urginea* spp., more often known as Indian squill, is a critically endangered medicinal plant that grows only in parts of India, Africa, and the Mediterranean. Magically, it has the ability to treat a wide range of illnesses afflicting humans, including cardiac disorders, cancer, jaundice, dropsy, asthma, epilepsy, dermatitis, and enuresis. It also changes the menstrual cycle and has abortifacient effects. As a pesticide, it may be used to kill rodents, insects, and fungi. Diverse genetic and chromosomal differences between *Urginea* populations were currently being investigated. Conserving this endangered species also requires a strong focus on biodiversity and germplasm collecting. There is still a lot to learn about this species, from the most fundamental taxonomic work to the most advanced molecular developmental investigations. In addition, it is a rich source of unexplored organic compounds that have the potential to serve as bioactive molecules. Research into genetic diversity and genomics is as popular as ever..

**Key words:** *Urginea*, Medicinal plant, Phytochemicals, biodiversity, botanical pesticides

### **Introduction**

German botanist Adolphe Steinhill (1834) identified 55 species and named the genus *Urginea* after the Arabian tribe Ben *Urginea*. This genus was classified by Lindley (1836) as belonging to the Scilleae tribe. *Urginea* is a fascinating and diverse genus that includes hundreds of species found in places like India, Africa, and the Mediterranean. Nine species belonging to the genus *Urginea* were the most frequent in the Indian context. Five species of this genus are native to India, according to a taxonomic immediately after the first shower followed by severe summer. The second category produces vegetative leaves along with the inflorescence axis soon after the first monsoon showers review by Deb & dasgupta<sup>13</sup>. The Indian

squill, or *Urginea*, is used in both pharmacology and agriculture. The ancient Egyptians found that it was effective against swelling, vomiting, and coughing. As an expectorant, it is still used in several over-the-counter cold remedies today. There is ongoing investigation on the efficacy of bulb extract as a cardiotonic employed by the ancient Romans. Despite being approved by the German commission in 1985 as 'E' for cardiac insufficiency, the use of the digitalis glycoside squill components is prohibited in the United States. More than a century of research has shown the efficacy of some squill varieties as rodenticides<sup>7,23</sup>. The species of *Urginea*, especially *U.*

*indica* is highly polymorphic with two distinct categories. The first category is very unique with underground bulbs producing inflorescence without vegetative leaves, immediately after the first shower followed by severe summer. The second category produces vegetative leaves along with the inflorescence axis. German botanist Adolphe Steinhill (1834) identified 55 species and named the genus *Urginea* after the Arabian tribe Ben *Urginea*. This genus was classified by Lindley (1836) as belonging to the Scilleae<sup>33</sup> tribe. *Urginea* is a fascinating and diverse genus that includes hundreds of species found in places like India, Africa, and the Mediterranean. Nine species belonging to the genus *Urginea*<sup>21</sup> were the most frequent in the Indian context. Five species of this genus are native to India, according to a taxonomic immediately after the first shower followed by severe summer. The second category produces vegetative leaves along with the inflorescence axis soon after the first monsoon showers. review by Deb & dasgupta<sup>13</sup>. The Indian squill, or *Urginea*, is used in both pharmacology and agriculture. The ancient Egyptians found that it was effective against swelling, vomiting, and coughing. As an expectorant, it is still used in several over-the-counter cold remedies today. There is ongoing investigation on the efficacy of bulb extract as a cardiotoxic employed by the ancient Romans. Despite being approved by the German commission in 1985 as 'E' for cardiac insufficiency, the use of the digitalis glycoside squill components is prohibited in the United States. More than a century of research

has shown the efficacy of some squill varieties as rodenticides<sup>7,23</sup>. The species of *Urginea*, especially *U. indica* is highly polymorphic with two distinct categories. The first category is very unique with underground bulbs producing inflorescence without vegetative leaves, immediately after the first shower followed by severe summer. The second category produces vegetative leaves along with the inflorescence axis

soon after the first monsoon showers.

#### **Classification**

Division	–	Liliophyta
Class	–	Liliopsida
Order	–	Liliales
Tribe	–	Scilleae
Family	–	Liliaceae
Genus	–	<i>Urginea</i>

The first known written reference to the squill dates back to the Egyptian Ebers Papyrus, which was written in 1500 B.C. It includes a traditional remedy for heart disease that was made by boiling together certain amounts of squill 30 years ago. There is documentation of sea squill's discovery and usefulness in ancient Greek texts. As an ancient Greek physician, botanist, and former pupil of Aristotle (371–287 B.C. ), Theophrastus writes on how effective it is in restoring human health. Diuretic, anti-jaundice, expectorant, anti-colic, and anti-asthmatic compounds were all given their own chapter in Dioscorides' (about 40–90 AD) Encyclopedia of Medical compounds. properties of sea squill with the detailed medical preparations and prescriptions.





Fig 2. Distribution- Populations of *Urginea indica* Kunth.



1. Kushahagar
2. Shimoga
3. Basavanahalli (Mysore)
4. Ranganghatta island (Mandya)
5. Chamundi Hills at Mysore
6. K.R.S at Mandya
7. Gopal Swayy betta, Chamaraingar District
8. Dopper Gowdara Para, Chamaraingar
9. Channamallipura (North), Gundlupet
10. Channamallipura (South) Gundlupet
11. Channamallipura (East) Chamaraingar Dist.
12. Channamallipura (West)
13. Maddur Forest Range – Mandya
14. Gotur – Hassan
15. Bangaswadi, Mandya
16. Sringeri (Kudremukh Forest Range), Chikkamagalur
17. Malahalli (Nanjangud)
18. Pityy village – Hiriyur – Chitradurga
19. Trichendur - Kerala
20. Anchipura – Gundlupet
21. Kalapura – Nanjangud
22. Tuticorin island – Tamilnadu Ramanathapuram Dist.
23. Handinuru - Mysore
24. Papanasini – Tirupathi – Andhra Pradesh
25. Gulbarga – Gulbarga
26. Karighatta – Srirangapatna – Mandya
27. Chamundi Hills – Mysore
28. Bellur – Tumkur Dist.
29. G.K.V.K. Campus (Bangalore)
30. Sholapur, Bombay
31. Nagamangala – Mandya Dist.
32. K.R.S. at Mandya
33. Kuvempu University (Shimoga- Bhdavathi)
34. East coast Kadarur

It also referred squill as vegetable or salad with a note aau, Basal El – Faar, BasalEl – Onsol, Basal

	Nsool, Skeletoura
Hebrew	Hatsav Matsu
French	<i>Scille maritima</i>
Italian	<i>Cipolla Marina</i>
Portugese	<i>Cila Maritime</i>
Arabic	Feraoun
Hindi	Jangli Kanda, Jangli dungli
Tamil	Narivengayam, Kaattuvengayam
Kannada	Kadu Irruli, Vana Palandu,
Kadu	Bellulli, Naaiyiulli, Seeme nari Eerulli, Shiru Naarangaddhe

**Synonyms** Basal tal ghansar, *Bulbo de escilla*, *Charybdis martima*, *Drimia maritime*, Ghansar, Meerzwiebel, Pharmacist'ssquill

**New Names** *Cebolla Albarrana*, *Drima indica*, *D. Maritima*, European squill, Mediterranean squill, *Scilla maritime*, White squill, Sea onion, Sea squill bulb, *Urginea scilla*, *Urginea indica*, *Scilla indica*

of caution about its poisonous nature. Pliny the Elder, the Roman author and naturalist (23 AD 79), categorized squill in to three types which included an edible one with a less pungent taste called 'Epimenides', describing the other two types by differentiating its appearance, taste and use as medicine. It was evident from fourteenth century documentations of Northern European countries about the use of squill vinegar prepared from dried squill. The Book of nature of Konrad of Megenberg, German scholar (1309 to 1374) referred a chapter on squill, describing it as mouse onion due to its rodenticide property. He identified edible squill varieties and also distinguished it as anti jaundice, anti dropsy and diuretic. Konrad of Megenberg also states that the Squill will cause abortion in pregnant women. The subject of the squill through history refer to the work of

J. Stannard (1974), in ancient and medieval *Materia Medica* with special reference to its application for treating dropsy<sup>53</sup>.

Language	Names
English	Sea Onion, Wild Onion, Indian squill, Red squill
Spanish	<i>Cebola Albarra</i> , <i>Cebola Chirle</i> , Esquila
Greek	Basal Fra

**Plant Description**

*Urginea indica* Kunth. commonly called as Indian Squill is a perennial geophyte with fibrous roots of six to ten inches of length, proceeding from the base of the bulb is a scapigerous herb. The rounded conical, pear shaped bulbs with white transparent outer scales are about the size of an big onion, consisting of fleshy coats which are thin and papery red or orange brown in colour enclosing each other completely. The phyllotaxy exhibited is whorled hysternanthus or synanthus. The bulb, which is usually three fourth immersed in the sand sends several long linear lanceolate, radical, cauline, lorate, sessile, pointed and undulated shining, dark green leaves with a base sheathing, becomes two feet when fully grown. From the middle of the leaves, around, smooth, long, terete, stiff and narrow succulent flower stem rises, one to three feet high terminating in a long, raceme, with close spike of whitish flowers, which stand on purplish peduncle. The flowers are bisexual, hypogynous, companulate bracteate and dropping. The flowers bloom in April and May after first shower followed by oblong capsules. The bracts are solitary, with long or short pedicel. The

perianth is lanceolate, subequal in two whorls of three each outspreading, free to the base or very near **Vegetative Phase**

one or few nerved at the centre. Stamens are six in number and are freely adherent to the base of the perianth segments. Anthers are ablong, dorsifixed, versatile in nature. Ovary is oblong or narrowly ovate, sessile superior, syncarpous, trilocular with short, thick and narrow style. It has subglobose stigma, with brownish capsule shaped shiny winged seeds, clustered, superposed and compressed in




reproductive phase features. The predetermined parameters of **Reproductive Phase** were assessed in two consecutive seasons to study and differentiate morphological variations from each other. At the population level the vegetative characters shown great variations on dependable taxonomic characters and mean time the reproductive characters shown insignificant uniform variations<sup>52,37</sup>.

#### Floral Biology

The studies on floral biology in *Urginea indica* shows that the species of the present study are self incompatible and cross pollination is mostly by insects. Pollen fertility is observed as 82% with no seedsetting. The data on the population studies have indicated that reproductive isolation through a difference in flowering period and blooming time as one of the factor that might have been played an important role in speciation and plausibly in the evolution of the genus.


#### Pollen Grains<sup>39</sup>

Pollen grains are monosulcate oblong with a single colpae extending from one end to the another. But a careful analysis of pollen grains in populations of *U.indica* revealed finer differences especially with regard to the size and the quality of exine reticulation.




**Anatomy** <sup>38,50</sup>

Sixteen populations of *Urginea indica* (Kunth). Liliaceae were examined to provide the first detailed description of leaf anatomy following the methods employed by Johansen. The populations were distinguished into two types based on the fleshy and watery leaves. Populations vary with the features like presence or absence of thick cuticle, larger or smaller areoles and clear vascular bundles. The mesophyll cells with intercellular spaces filled with heavy and moderate wax deposition. Larger and smaller epidermal cells, palisade like tissue these characters along with other parameters plays an important role in delimiting the populations. Idioblastic cells containing raphide bundles with calcium oxalate crystals occur in the lower side of the mesophyll. These variations between populations are of taxonomic significance



#### Scanning electron microscopic studies of leaf surface



#### Leaf cuticular ornamentations stomatal variations and the differences in wax deposition, presence of raphides etc play an important role in delimiting the populations

of *Urginea indica*. It is suggested that leaf surface characters can be used as secondary or supporting character in biosystema studies. Shiva Kameshwari et al. (2001)


#### Stomata

Studies were conducted on epidermal micromorphology of twenty one different populations of *U.indica* collected from various localities of


Karnataka. They showed the presence both amphistomatic and amocytic type of stomata on the upper and lower epidermis as reported as contributing variations in stomatal index. The comparative study on the matured leaves of different populations showed less variation in stomatal frequency, area of stomatal aperture and the size of guard cells<sup>38</sup>.

### Defense Role

Plant defend<sup>1</sup> themselves against attack from herbivores has been the subject of considerable interest over many decades. Plant structural traits such as raphides play an important role in protecting plants from herbivore attack. These raphides are evolved as a result of the response caused by the other environmental stimuli on the mesophyll intercellular spaces idioblastic cells containing raphide bundles and different phenotypes of crystalloid inclusions embedded in poly saccharides. Such crystals usually have backward oriented surface barbs capable of increasing damage to the mouth of grazing animals. Raphides are also responsible for producing mild inflammation and itching<sup>12</sup> when rubbed on the skin. Therefore, raphides take part in both mechanical and chemical irritation when they come into contact with tender tissues of soil living worms and herbivores<sup>11</sup>.



The roots of *U. indica* play an important role in storing and utilizing water and nutrients thus protecting the plant from drought stress and environmental hazards. Raphides are present in all organs mean while bulbs showed richness in raphides. Thus calcium oxalate crystals occur in different forms in *U.indica* and perform various functions including herbivorous defense<sup>25</sup> calcium regulation (Volk et al., 2002) and are associated with



### Distribution And Ecology

Sea squills with survive in areas with a little as 100 mm annual rainfall but are restricted to coastal regions. Quoting from a study by Kamal Hassan et al (1970) *U. maritima* is a polymorphic species with different varieties and forms Squill growing in Egypt show three distinctive features regarding the morphology of the bulbs. One with moderate size and reddish tinge, the second with white tunics and small size while the third with dark red tunics and very large bulbs<sup>28</sup>. This morphological variation is directly tied to soil type. In Egypt red bulbs inhabit soils of sand stone origin. Whereas white squill bulbs are found in soils of limestone origin. Whereas *U. indica* endemic to India, Africa and Mediterranean regions is found in a wide habitats ranging from desert, shrub, grassland, dunes and forests, soil conditions in which the squill is found are equally varied including sand, clay, calcareous, acidic and saline<sup>37</sup>. Recently it is found growing in platinum rich area Shiva Kameshwari and Paramasivam (2011) yet to be published

### Germplasm

India being one of the natural centre of origin of *Urginea Indica* with large germplasm base, it is gaining immense global importance in view of its potential for multiple uses. Due to this, there is great scope & need to collect all species & their population & conserving them in one place. Therefore, there is immediate need to devise a program aiming at systematic collection documentation & characterization of *Urginea* geruplasm in India. *Urginea* collections in India. And also a detailed germplasm catalogue, comprising passport information on an the important attribute of different *Urginea* species including molecular ID cards is in progress.

refractory to phenobarbital and phenytoin without causing respiratory depression<sup>5</sup>. Tartronic acid is used as an oxygen scavenger<sup>3</sup> in United States. Acid group shows the presence of mindereru's spirit which initiates

Mediterranean region, cultivated for drug extraction at Mediterranean regions, southern states of American and in the Caucasus regions. Turkey squill is harvested

HO OH structures of some Bioactive Compounds

and traded locally from the wild area of different countries around the Mediterranean sea. Squill grows slowly and the bulb is not ready for harvesting until the sixth year when it produces its first lower stem. If it is allowed to continue growth then for several years the flowers can be harvested as cut flowers (Economic Botany). In India strict control of squill collection needs to be maintained in order to protect its wild populations. It is in progress in few hilly regions. For Eg: Siddarbhata, Tumkur and Coimbatore, Wellingiri hills. Vegetative propagation was found to be stronger in desert populations of *U. undulata* compared with the Mediterranean populations of *U. maritima*.

#### Phytochemicals

Squill contains a large number of related steroidal cardioactive glycosides. Scillaren A and proscillaridin A, are the major glycosides found in the bulb in greatest concentration<sup>59</sup>. Other constituents found in squill include flavonoids, carbohydrates, antifungal glycoproteins, steroids, alkaloids, esters and saponins. Main active ingredients of squill are

steroidal glycosides. Many new natural compounds have been isolated from *Urginea indica* by recent researchers are yet to be included in the organic chemicals repository<sup>23</sup>.

The phytochemicals extracted from the bulbs were found to be potentially bioactive. 2, 3-Butanediol is used as a cardiac stimulant<sup>8</sup>. Along with these other compounds such as paraldehyde, tartronic acid, quercetin, and mindereru's spirit were also identified. Paraldehyde – a polymer of acetaldehyde is used as a sedative, hypnotic and anti seizure. It is a potent anticonvulsant capable of controlling seizures. Steroids (corticosteroids) present in the bulbs were

used to treat psoriasis by indigenous people.

Quercetin, a potential bioactive molecule associated with wild onions were found significant in reducing the blood pressure by an average of five millimetres of mercury. Bufadienolides

were identified in different chromosomal races of

Indian squill *U. indica* (Shiva Kameshwari and Muniyamma, 2000). Identification of a novel 29 kDa glycoprotein with antifungal activity from Indian squill and its role in biological control were also being researched<sup>31,35,46</sup>.

**Table (1): The Important Compounds and the attached basic principal components responsible for bioactivity**

Sl.No	Activity	Compounds/ Basic Principal
1.	Acetylcholinergic	Glucose
2.	Anticarcinomic	Scillarenin
3.	Anticystic	Mannose
4.	Antidiabetic	Xylose
5.	Antiedemic	Glucose
6.	Antihepatotoxic	Glucose
7.	Antiketotic	Glucose
8.	Antirhinoviral	Scillarenin
9.	Antivaricose	Glucose
10.	Antiviral	Scillarenin
11.	Cancer preventive	Mucilage
12.	Cardiac	Scilliglaucoside
13.	Cardiotonic	Scillarenin
14.	Demulcent	Mucilage
15.	Diagnostic	Xylose
16.	Dye	Xylose
17.	Hypereglycemic	Glucose
18.	Memory enhancer	Glucose
19.	Pesticide	Scillarenin

The leaf flavonoids in different populations of *Urginea indica* showed variations in the possession of cyanidin, petunidin & pelargonidin, they differ in their Rf values. Few populations are dissimilar since they are characterized by the absence of one of the three above mentioned compounds<sup>54</sup>

#### Pharmacology and Medicinal Uses<sup>22</sup>

Squill glycosides were identified for its cardiotonic properties similar to digitalis from Pharaonic times. However, squill components are less potent than digitalis. Preparations for oral administration are enteric coated to prevent degradation from gastric acid. Meproscillaren, a semisynthetic derivative of proscillaridin, is absorbed orally and may be effective in some patients. Based on British Pharmacopoeias assay for digitals there is no differences between extracts of *U. maritima* and *U. indica*.

Squill induces vomiting. Vomiting may be preceded by a generalized increase in the flow of secretions, and therefore these compounds appear to exert an expectorant effect in sub-emetic doses. Methanolic extract of red squill have been used as hair tonics in treating Seborrhea and dandruff<sup>32</sup>. Red squill is not used medicinally but used as rodenticides. Squill has

been used traditionally as a cancer remedy and siliquoside, has shown activity in an experimental, cancer cell line<sup>16</sup>.

Squill is used in human homeopathy and phytotherapy and in veterinary science. It is administered orally, typically as a diuretic and functions by increasing blood flow through the kidney, emetic expectorant. In Greece fresh bulbs are distilled for medicinal use. The German commission E. Monographs suggests squill can be used for milder cases of heart insufficiency and also for diminished kidney capacity [European agency(1999), Blumenthal et al(1998)]. Sinistrin, an inulin like substance, is extracted from squill for use as a marker in diagnosis of renal problems.

Muscle pain from disease such as fibromyalgia, over work of a muscle which leads to cramps and contractures, contractures are continuous muscle contractions with associated chronic nagging pain. This type of chronic pain is difficult to treat. Use of squill extract for muscle pain has proved effective and developed as an analgesic<sup>59</sup>.

Psoriasis an inflammatory disease of the skin, where treatments are not available in Allopathic medicine but *Urginea indica* preparations are known to be used traditionally by many tribes and aborigines against psoriasis and many dermatological diseases.

#### **Cancer Activity**

The activity studies conducted *in vivo* and *in vitro* against mouse mammary carcinoma cells proved that purified 29 kDa glycoprotein from squill as anticancerous. This protein assayed against cloned line of human colon adino carcinomatic GC<sub>3</sub>/C1<sup>7</sup>, KB Ch<sup>f</sup>- 8-5 and KB-3-1 strain, using DMEM and RPMI media were also proved as anti cancerous.

#### **Insecticidal Activity**

The phytobiocative compounds extracted from *Urginea maritima* showed excellent mosquito repellent action<sup>34</sup>. Recent studies on *Urginea indica* remarkably shown its larvicidal action against *Aedes* larvae causing dengue fever. The 400µl of aqueous lyophilised extract showed cent percent mortality of the larvae within fifteen hours.

#### **Toxicology and Agricultural Uses<sup>22</sup>**

The Egyptian Alexandria plague epidemic of 1946 – 47, squill was used as rodenticide bait to control the spread of plague. Ab etel Gawad (1955) developed a Rat bait recipe using 1:3:1 proportion of white squill: flour: tallow with reasonable amount of salt by parted as one in three hundred portions, illustrates the potential toxicity caused by squill. Among the active principals, scillirosides shown greater bioactivity against rats exploiting the inability of rats to vomit the poison<sup>20</sup>. Because


squill-laced bait is vomited by domestic animals before a lethal dose can be absorbed, often it is considered to be a rat-specific agent.

Christos Georgiades book 'Flowers of Cyprus' portrays about the potential possibilities of squill components as insecticides. Against storage pests of barley and wheat farmers normally use sea squill bulbs.

The larval growth retardant as well as adult fertility depressant actions of bufadienolids compounds extracted from bulbs of sea squill were found effective against storage pests like red flour beetle, *Tribolium castaneum*<sup>34</sup>. Recent research about the squill extracts leading to decrease the load of toxicants as synthetic pesticides and drugs or Neti mert (2008) Over dosage affects pregnant women and also plays its toxic role in menstrual cycle. The bulb extract showed hypoglycemic activity. The alcoholic extract is found active against *Entamoeba histolytica* strain (C.P.Khare 2004) Rabbits, however, were found dead after they had chewed on fresh bulbs in the ground. Red squill a fine raticide, red squill plants should find a permanent home in the south west as a new specialty crop that should prove profitable to some farmers, increase the health of city dwellers and provide chicken farmers with better control of wasteful and pestiferous rats.

The studies on inhibitors in *U.indica* indicate their presence in leaves, bulbs and in seeds the concentration of the inhibitory substances in the bulb varies during its different growth phase<sup>29</sup>.

#### **Cytogenetics<sup>36,40,41</sup>**



Recent attempts were made by researchers to distinguish *Urginea sps.* based on chromosome



number and distribution to form an exclusive database through cytogenetic approach.

The general chromosome morphology of *U. indica* as a whole is quite distinct, the chromosome size ranges from very long to very short and often there is size difference in thirty two different population studies.

**Chromosome number variability of *Urginea***

No	Species	Chromosome
1.	<i>U. altissima</i>	2n = 20
2.	<i>U. aurantiaca</i>	2n = 21
3.	<i>U. burkei</i>	2n = 20
4.	<i>U. coromandeliana</i>	2n = 20
5.	<i>U. depreesa</i>	2n = 40
6.	<i>U. epigea</i>	2n = 32
7.	<i>U. fugax</i>	2n = 22
8.	<i>U. govindappae</i>	2n = 20
9.	<i>U. indica</i>	2n = 20
10.	<i>U. langii</i>	2n = 20
11.	<i>U. lydenburgensis</i>	2n = 32
12.	<i>U. maritime</i>	2n = 20
13.	<i>U. mouretic</i>	2n = 54
14.	<i>U. nigritiana</i>	2n = 60
15.	<i>U. polyphylla</i>	2n = 20
16.	<i>U. pretoreinsis</i>	2n = 20
17.	<i>U. rubella</i>	2n = 42
18.	<i>U. tenella</i>	2n = 20
19.	<i>U. undulata</i>	2n = 20

Cytological studies on populations of *U. indica* revealed the diploid, triploid, tetraploid, aneuploid and hexaploid nature of the populations through karyotype attributes using different parameters. (Shiva Kameshwari 1999)

Each cytotype studies differed distinctly in vegetative and floral characters. Consequently the thirty two cytotypes studied were observed by having distinct somatic complement of chromosomes. Thus, it lead into the recognition of different cytotypes among the natural populations. Several aneuploids have also been recorded in the recent investigations.

Polysomaty has been found to be a regular feature in plants which reproduce through vegetative means (Sen, 1973). The origin of such nuclei with varying chromosome numbers may involve various cytological mechanisms such as endomitosis, non disjunctions and duplication of chromosomes leading to polyploidy and aneuploid cells (Shiva Kameshwari and Muniyamma, 1999).

A team of researchers were still working on developing a cytological database on *Urginea indica* (Raghavan 1935).<sup>41,42,46,47,51,52,61</sup> Occurrence of cytomixis and its importance in evolutionary diversification of species were reported in diverse


species of angiosperms (Shiva Kameshwari and Muniyamma, 2001, 2008). Meiotic irregularities & variations in the chromosomal behaviour indicates that the populations could eventually be treated as chromosome races.

B-chromosomes are also recorded during mitosis ranging from one to ten for *Urginea Indica*.

**Karyotype**

The karyotype in *Urginea indica* is asymmetrical with a graded series in which ST and SM type chromosomes predominate. It further suggests that the numerical evolution by polyploidy and structural changes leading to intra karyotypic size differences of chromosomes and shifting of centromeres from median to submedian and sub telocentric have been concomitantly operating in the genus.

**Embryological studies**



**Microsporogenesis**

The anther is four lobed with two middle layers. The endothecium shows evidences of disintegration as fibrous thickening in the inner middle layer. The tapetum is granular with binucleate cells. Pollen grains are two celled. Microspore mother cells undergo reduction division resulting in the formation of isobilateral tetrads. Pollen grains show a smaller generative cell and a larger vegetative cell. Pollen grains shows smooth, exine and thin intine.

**Megasporogenesis**

Megasporogenesis and the female gametophyte development conform the monosporic eight nucleate polygonum type of embryo sac development. However, the organized mature Embryo sac has a broader micropylar part and a narrower chalazal region in *Urginea indica*.

**Plant Tissue Culture**

Three chromosomal races (diploid, triploid and tetraploid) of Indian squill were screened for the production of bufadienolide proscillaridin-A (PsA) and scillaren-A (ScA) in tissue cultures. Bulbs and inflorescence segments were cultured on Murashige-Skoog medium supplemented with various combinations of plant growth factors, and /or coconut milk and yeast extract. Callus formation was induced from bulbs of diploid and triploid genotypes and from


inflorescence segments of tetraploids. The shoot buds developed into small bulbous plantlets in plant PGF-free medium. Somatic embryogenesis was observed in the long-term callus cultures. Shoot differentiating calli (callus with 10-20 shoot buds/8-12 weeks old) contains low levels of PsA in both the diploid, triploid and tetraploid races. All regenerated bulbous plants produced both PsA and ScA Jha (1991)<sup>27</sup>.

**Molecular Biology and Genetics**

Molecular analysis primarily depends on the availability of high purity DNA sample and reproducible protocols for employed marker analysis. Here we present a first report on the optimization of DNA isolation and PCR conditions from bulb tissues of *Urginea indica*. The bulb tissues were used to study due to non-availability of leaf material during all the seasons. However, isolation of DNA from the storage tissue like bulb was particularly, challenging because of their high levels of polysaccharide and protein contents.

Existi


ng



resulted in pure genomic DNA that was amenable to restriction digestion and PCR amplification after protein and polysaccharide contamination had been removed. When compared to reference DNA samples, the extracted DNA displayed a signature restriction digestion pattern using four enzymes: Alul, Hind III, ECO RI, and BAM H. The first standardized procedure for DNA isolation and PCR - RAPD analysis in *Urginea indica* has opened the door to further in-depth molecular studies of this vital medicinal plant.


*Urginea indica*'s significant therapeutic potential makes it a promising option for future medical uses. Comparative molecular characterization of distinct plant species and populations may reveal the genetic basis of variances in plant qualities, paving the way for further research into and development of these features. For this reason, it is possible to create marker systems like RAPD, RFLP, SSR, AFLP, etc. Furthermore, the study of evolutionary relationships between species is a valuable resource. There is, therefore, an urgent need to conduct a comprehensive molecular characterisation of *Urginea* species in India<sup>15,54</sup>.

**Biochemical Studies**  
29kDa protein



**Antifungal Activity**

protocols described for other liliaceous species employed non bulbous materials; hence, were in effective for the present study. Therefore, several modifications



*Fusarium oxysporum*

*Rhizoctonia Solanii*


Purification by acid precipitation, Diol-Gel filtering, and C18 reverse phase HPLC yielded a homogeneous antifungal protein from *Urginea indica* bulbs. Periodic acid-Schiff (PAS) staining and a calculated molecular weight of 29 kDa confirmed that the detected antifungal molecule is a glycoprotein. It seems that the glycan portion of the molecule is involved in anti fungal action, since this property is lost during periodic oxidation of the 29 kDa glycoprotein. The isolated glycan-producing protein has an N-terminal amino acid sequence of SQLKAXIXDF and showed no sequence resemblance to known antifungal proteins.

Immunolocalization studies made use of a polyclonal anti serum that had been produced against a purified protein. The protein in *U. indica* has antifungal properties. It entirely stops *Fusarium oxysporum* from producing both spores and hyphae. In an in vitro experiment, the Mr 29kDa protein found in the purified formulation was shown to actively inhibit the development of the fungal diseases *Fusarium oxysporum* and *Rhizoctonia solani*. The Mr 29kDa protein, whose chitinase function was deduced from its amino acid sequence, has striking homology to the family 17 glycoside hydrolases.

The chitinase from *Urginea indica* has a conserved pattern typical of the signature 1 of family 19 glycoside hydrolases but was missing the cysteine-rich N-terminal domain seen in class I chitinases. It was a typical class II chitinase from family 19, sharing 70% sequence similarity with the *Hordeum vulgare* endochitinase (26kDa). The Mr 29kDa *U. indica* protein is likely a class II chitinase due to its molecular weight, absence of an N-terminal cysteine rich region, and strong similarity to the *H. vulgare* endochitinase. The chitinolytic activity of the Mr 29kDa protein is likely responsible for the antifungal effect.

#### Chitinase Protein Research on *Urginea indica*

Chitinase (EC 3.2.1.14), which is found in barley, wheat, rye, maize, garlic, etc., has a particular hit in the protein sequence below, indicating that it belongs to the lysozyme-like superfamily. The sequence was also found to be completely unique, sharing no similarities with any other known fungal proteins.



The FASTA file sequence is given below:

```
-----
--- NH3-
SVSSIVSRAQAQPPKPSHAFDRMLLHRNDGAC
Q
AKGFYTYDAFVAAAAAFSGFGTTGSADVQKRE
L
AQTSHETTGGWATAPDGAFWGYCFKQER
GAS
SDYCTPSAQWPCAPGKRYGRGPIQLSHNYN
YG
PAGRAIGVDLLANPDLVATDATVSADRAAG
RVPFGFVITNIINGGIECGHGQDS
-COOH
-----
```

#### Percentage Contents of *Urginea indica*

SI No	Contents	Percent/100gms
1	Protein	8.36
2	Carbohydrate	66.25
3	Fats	0.32
4	Fibre	12.30
5	Moisture	6.16
6	Phosphorus	0.13
7	Acid value	7.23
8	Total ash	6.61
9	PH 5% solution	4.68
10	Fat	0.32
11	Energy	301.32 Kcal.
12	Calorific Value	3503 cal./gm

#### Microbiology

Indian squill *Urginea indica* is showing antifungal activity against *Fusarium oxysporum* which effects Nanjangud Rasabale, to *Sclerotium rolfsi*, *Alternaria tenuissima* and *Rhizoctonia solanii*

which kills nursery plants<sup>11</sup>.The extract was prepared with Methanol and are subjected for preliminary phytochemical and physicochemical analysis. The total ash content, acid insoluble and water soluble ash content were evaluated

along with the fluorescence characteristics of the methanolic extract of wild onion sps. The presence of primary and secondary metabolites such as *carbohydrates, proteins, alkaloids, phenolic compounds, saponins* were confirmed through preliminary phyto-chemical analysis.

The extract was found to possess anti-bacterial activity in *E. coli, S. aureus and P. aeruginosa* isolated from infected patients. The Minimum inhibitory concentration (MIC) was also evaluated by 'Tube dilution' method and the result was found to be considerably effective against selected pathogenic bacteria. Such an effect might contribute in explaining the traditional use of wild onion sps, *Urginea indica* in the treatment of wound healing.

The antioxidant activity was estimated by using DPPH free radical scavenging assay and the activity was increased with increase in concentration of methanolic fraction of wild Onion sps. The fractions of wild onion sps are free radical scavengers and are able to react with the DPPH radical, which might be attributed to their electron donating ability. Thus suggested the antioxidant components in this Wild Onion sps capable of reducing oxidants and scavenging free radicals. This also indicates that, tubers of wild onion, *Urginea indica*' are of therapeutic potential due to their high free radical scavenging activity. The presence of high amount of saponins justifies the practice of treatment for disturbances in the gastrointestinal tract by traditional healers.

The role of phyto-chemical constituents of this Wild Onion, *Urginea indica* sps in traditional medicine treatment was discussed. Hence, the formulation of extract of *Urginea indica* needs to be purified using biophysical techniques towards development of a potential drug/ lead molecule against microbial infection, inflammation and wound healing respectively.

#### Heavy Metal Analysis

No	Metals	Amount (mg/100gm)
1.	Silica	290.00
2.	Alumina	585.00
3	Iron	230.00
4.	Calcium	80.00
5.	Magnesium	22.00

6.	Sodium	65.00
7.	Potassium	120.00
8.	Copper	0.16
9.	Manganese	0.14
10.	Zinc	0.38

#### Conclusion

*Urginea indica* commonly called Indian squill is considered to have medicinal value and is largely used as an expectorant, cardiac stimulant, in treating rheumatism, dropsy, edema, gout, asthma and as an anticancer agent. Due to these properties, the squill bulbs have found place in British and European Pharmacopoeias. Squill bulbs have long been used as a source of natural product with Pharmaceutical and biocidal (rodenticide, insecticides and fungicide) application<sup>34</sup>. Due to unawareness we have lost many populations of *Urginea indica* and it is necessary to initiate awareness, conservation and cultivation of *U. indica*. Anthropogenic pressures such as habitat degradation are largely responsible for genetic depletion and loss of genetic diversity. New means and approaches are to be worked out for germplasm conservation and sustainable utilization of this economically important medicinal plant. The standardization of agro techniques and propagation program is in progress. Population studies have also been made in *U.indica*. An attempt has been made to enquire into the morphological variations which lead to evolutionary divergence of populations of *Urginea indica* Kunth. In particular, there were considerable morphological variations within the species. The thirtytwo cytotypes showed distinct morphological differences, in shape, size and colour of bulb and leaves, the length of inflorescence, and flower colour. The morphological complexity is accompanied by high degree of cytological variations. Preliminary measurement of reproductive characters have shown that no noteworthy results may be obtained in this characters except pedicel length but the vegetative character deviate significantly. The flowering and blooming time varied also played an important role in differentiating the populations. These morphological differences, have a genetic basis and would be worthy in recognizing them as a separate sub specific taxon.

Morphological and Cytological variations, revealed the presence of diploid, triploid, tetraploid, aneuploid and hexaploid populations and these were explored for their karyotype attributes using different parameters. We also report the presence of polygony of eight celled embryosac development as well as *in vitro* culture and regeneration in one

population. The bioactive principles of *U.indica* were extracted and tested for their antifungal and anti-cancer activities. The findings of the study indicated that the crude bulb extract can be used for various purposes. In the present study, we have isolated and studied the antifungal activity of a 29 kDa protein against, *Fusarium oxysporum* and *Rhizoctonia solani*. We have also identified the antibodies developed against the protein which neutralizes its activity as fungicide. N-terminal amino acid sequence analysis showed that the purified antifungal compound protein is found to share high homology to 29 kDa Endochitinase group. *Hordeum vulgare* with a conserved domain Glyco hydro 19, characteristics of Chitinase Class I proteins. These data suggest that the purified antifungal compound is a Putative Endochitinase and could be a bonafide member of class II Chitinase showing anticancer activity. This investigation is aimed at making popular Indian squill an economic and medicinally important plant for India<sup>35</sup>

### Acknowledgement

The valuable suggestions and support from Prof. Sumitha Jha, Centre for Advanced studies, Department of Botany, University of Calcutta, is well appreciated in the preparation of this manuscript. We are very much thankful to Dr. Palani G. Periasami, Chairman, and Mrs. Vishalakshi Periasami, Vice Chairman, PGP Educational Welfare Society, Chennai, Tamil Nadu for providing the required environment for making this review article.

### References

1. Ab etal Gawad Hussein. (1955). Changes in the epidemiology of plague in Egypt, 189 – Bull World Health Organ. (1955). 13(1): 27 –48, PMID; PMC2538042.
2. Agarwal A.A., Fishbein M.(2006). Plant defence syndromes Ecology 87:S132,S149
3. Ahmed, S., Ellis, J.C., Kamwendo, H. (2006)Efficacy & Safety of intranasal lorazepam versus intramuscular paraldehyde for protracted convulsions, in children ; an open tandomised, trial Lancet 2006 : 367: 1591-97.
4. Airy Shaw H.K., (1966). A Dictionary of flowering plants and freus 8<sup>th</sup> Edition (Willis J.S.Ed. Cambridge university press Cambridge).
5. Armstrong D.L.,BattiuM.R.,(2001). pervasive seizures causes by hypoxie
6. Bizot paul M.,Bruee & Bailey., Peter D.Hicks use of tartronic acid an oxygen scavenger patent number 5,750,037 May12 1(996).
7. Blumenthal M, Gruewald J, Hall T, Riggins, C., Rister R.(1998). The complete German Commission E. Monographs: Therapeutic Guide to Herbal Medicines. Austin, TX: American Botanical Society.Bridgitte Kopp.,Krenn L., Draxler M.,(1996) Bufadienolides from *Urginea maritima* from Egypt. *Phytochemistry*, **42**: 513 – 522.
8. Changes in the epidemiology of plague in Egypt, 189 – Ab etal Gawad Hussein. (1955). Bull World Health Organ. (1955). **13**(1): 27 – 48, PMID; PMC2538042.
9. Christos Gerogiades, Nicosia, (1987). flower of Cyprus, plants of medicine. ISBN, 9963 – 7540 – 1-5.
10. Civelek H.S., Weintraub P.G., (2004). Efect of two plant extract on larval leaf neiver *Liriomyza trifolii* (Diptera:Agromyzidae) tomatoes. *Jour of Economic Entomology* **97**:1581-1586.
11. Cogne A.L., Marstou A., Mevi S.,Hostettmann K., 2001. Study of two plantsused in traditional medicine in zimbabwe for skin problems and rheumatism *Dioscora sylvatica* & *urginea altissima* *Jour of Enthopharmacology* **75**:51-53
12. Deb DB, Dasgupta S, (1987). On the identity of three species of *Urginea* Liliaceae J.Boulay *Nat.Hist.Sci* **84**:409-412.
13. Deepak A.V., Thippeswamy G., Shivakameshwari, M.N., Salimater B.P., (2003). Isolation and characterization of a 29 kDa glycoprotein with antifungal activityfrom bulbs of *Urginea indica*. *Biochem Biophys Res. Comm.* **311**: 735 – 742.
14. Doyle J.J.,Doyle J.L., (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue .*Phytochemistry Bull* b:11- 15.
15. Duke J.A., (1985). C.R.C. Handbook of Medicinal Herbs, Boca Raton, FL: CRCPress:
16. *Economic Botany Volume*, **41**, Number 2, 267 – 282, DOI: 10.1007/BF02858974 Red squill (*Urginea Maritima*, Liliaceae) H.S. Gentry, A.J., Verbiscar and T.F. Banigan Red Squill.
17. European Union health files, 1999.
18. Franceschi, V.R., Nakata, P.A. (2005),Calcium oxalate in plants ; formation & function. Annual

review of plant biology 56: 41-71.

19. George Fawaz & Hildegard (1953) Rat poison from white squill. Meyer Br. J. Pharmacol Chemother, 1953, 8(4) 440-443, PMID ; PMC 1509361.
20. Hemadri K., Swahari S., (1982). *Urginea nagarjunae*, A new species of Liliaceae from India. *Ancient Sci. Life* 105-110
21. <http://www.drugs.com/npp/squill.html>.
22. <http://www.emedicinehealth.com/squill/vitamins-supplements.html>
23. [http://www.integrativepractitioner.com/article\\_ektid14976.a](http://www.integrativepractitioner.com/article_ektid14976.a)
24. Hudgins JW., Kreckling T., Franceschi V.R., (2003). Distribution of calcium oxalate crystals in the couifers a constitute defense mechanisms, *New phytologist* **159**:677-690.
25. Jha S., Sen S., (1981), Bufadienolides in different chromosomal races of Indian squill. *Phytochemistry*, **20**: 524 – 526.
26. Jha.S, Sahu-N-P, Mahato S.B., 1991. *Urginea indica* bulb culture, inflorescence culture medium optimization, effect of genotype on callus culture induction, organogenesis, somatic embryogenesis, propagation of bufadienolide prep. i.e, PsA Prep., ScA prep. *J. Plant-cell-Tissue-Organ-Culture*: **25**, 2, 85-90
27. Kamal Hassan Batanotny and Taha Khalifa, (1970). Contribution to the autecology of *Urginea maritima* in Egypt,
28. Khare C.P., (2004). Encyclopedia of Indian Medicinal plants springer verlag Berlin Heidelberg New York IS Br **3**-540-20033-9 464-465
29. Kreig, M.B., (1966). Green Medicine. The search for plant that heal, 4<sup>th</sup> print. Chicago, New York, San Francisco.
30. Krenn, L, Jelovina M, Kopp B. (2000). New Bufadienolides from *Urginea maritima sensu strictu*. *Fitoterapia*. **71**: 126 – 129.
31. Leung A.Y. (1980). Encyclopedia of common natural ingredients used in Food, Drugs and Cosmetics. New York, NY: *J. Wiley and Sons*.
32. Lindley J., (1836). Natural system of Botany. 353-354.
33. Pascual – Villalobos, M.J. (2002). Anti insect activity of bufadienolides from . P. 564 – 566, In. J. Janick and A. Whipkey (eds). *Trends in new crops and new uses*. ASHS Press, Alexandria, VA Anti-insect activity of bufadienolides from *Urginea maritima*.
34. Sandya 7 Shenoy., Shiva kameshwari M.N., Swaminathan., Gupta M.N., (2006). Major antifungal activity from the bulbs of Indian quill *Urginea indica* is a chitinase. *Biotechnology progress* **22**(3) 631-637
35. Sen.s., (1973). Polysomates and its significance in Liliales *cytologia* **38**:737-757 Shiva Kameshwari 1995, Biosystematic studies in some members of liliaceae, Ph.D. Thesis University of Mysore.
36. Shiva kameshwari M.N., (2011). Epidermal Micromorphology in populations of *Urginea indica* Kunth. Liliaceae. *International jour of Engineering Sci & technology*. **3**:3816-3824.
37. Shiva kameshwari M.N., (2011). pollen morphology in some members of Liliaceae. *International jour of Engineering Sci & technology*. **3**:3825-3830.
38. Shiva Kameshwari, M.N. and Muniyamma, M. (2009). Cytomixis in *Urginea indica* Kunth. Liliaceae, *J. Swamy Bot. Cl.* **25**: 47 – 50.
39. Shiva Kameshwari, M.N. and Muniyamma, M., (1999). Cytological studies in *Aneuploid Indian squill*. *Cell and Chromo Res.* **21**(2): 53 – 57.
40. Shiva Kameshwari, M.N. and Muniyamma, M., (1992). Karyomorphological studies in *Urginea indica* Kunth. Liliaceae. *J. Mysore Univ.* **32**: 518 – 522.
41. Shiva Kameshwari, M.N. and Muniyamma, M., (1999). Karyological studies in *Urginea polyphylla* Hook. F. Liliaceae, *J. Swamy Bot. Cl.* **16**: 91 – 93.
42. Shiva Kameshwari, M.N. and Muniyamma, M., (1999). Karyological studies in *Scilla indica* Baker. Liliaceae. *J. Swamy Bot. Cl.*, **16**: 87 – 90.
43. Shiva Kameshwari, M.N. and Muniyamma, M., (1999). Karyomorphological studies in *Drimiopsis kirkii* Baker. *Cell and Chromo Res.* **21**(1):

- 32 – 37.
44. Shiva Kameshwari, M.N. and Muniyamma, M., (2000). Cytological and chemical examination of Indian Squill. Book.
  45. Shiva Kameshwari, M.N. and Muniyamma, M., (2001). Cytological studies in *Scilla indica* Baker., *J. Swamy Bot. Cl.* **18**: 41 – 44.
  46. Shiva Kameshwari, M.N. and Muniyamma, M., (2001). Cytology of *Sansevieriacylindrical* Bojer. Liliaceae. *J. Swamy Bot. Cl.* **18**: 45 – 48.
  47. Shiva Kameshwari, M.N. and Muniyamma, M., (2001). Cytomorphological studies in three species of the genus *Chlorophytum* Ker Gawl, Liliaceae. *Taiwania* **46**(4): 307 – 317.
  48. Shiva Kameshwari, M.N. and Muniyamma, M., (2001). Scanning Electron Microscopic studies on leaf surface and seed surface in some taxa of Liliaceae. *Phytomorphology.* **51**(2): 137 – 144.
  49. Shiva Kameshwari, M.N. and Muniyamma, M., (2004). Karyomorphological studies in 14 populations of *Urginea indica*. Kunth. Liliaceae. *Beitr. Biol. Pflanzen.* **73**: 377 – 394.
  50. Shiva Kameshwari, M.N., Thara Sarawathi, K.J. and Muniyamma, M., (2010). Morphological Variations in population of *Urginea indica*, Kunth. Liliaceae. *J. Natural and Applied Science*, **2**(2): 280 – 289.
  51. Squill in a ancient and medieval materia medica, with special reference to its employment for dropsy. *J. Stannard Bull N.Y. Acad Med.* (1974). June, **50** (6): 684 – 713, PMID: PMC1749406.
  52. SS.Harini, M.Leelambika, et al.(2008). IJIB. Optimization of DNA isolation and PCR RAPD methods for molecular analysis of *Urginea indica*. vol **2** no.2, 138
  53. Steinhill.A.,1834.Note nur le genre *Urginea* Nouvelleent forme dans la familie de Liliaes- *Ann.Sci.Nat.II Bot.***1**:321-332.The European Agency for the Evaluation of Medicinal Product, (1999). Summary report *Urginea Maritima*.
  54. The Medusa Database (<http://medusa.maich.gr>) and references contained there in.
  55. The pharmacological Action of Digitalis, Strophanthus, and Squill on the Heart G.S. Haynes *Biochem. J.* (1906). **1**(2): 62 – 87, PMID: PMC1276124.
  56. Vahdettin Bayazit, Vahit Konara Analgesic Effects of Scilliroside, Proscillaridin – A and Taxifolin from Squill Bulb (*Urginea maritima*) (2010) on Pains Digest *Journal of Nanomaterials and Biostructures* Vol. **5**, No. 2. P. 457 – 465, Analgesic Effects of Squill.
  57. Volk GM, Lynch-Holm V.M., Kostma T.A., Gess L.J., Franceschi, V.R. (2002), The role of druse & raphide calcium oxalate crystals in tissue calcium regulation in pistia stratiotesleaves plant biology 4: 34-45.
  58. Yadav, S.R., Dixit G.R., (1990). Cytotaxonomical studies in Indian *Urginea* stein hill, species.*cytologia* **55**:293-300