




ISSN 2395-650X

International Journal of
Life Sciences Biotechnology Pharma Sciences

IJLBPS



www.ijlbps.org

E-mail: editorijlbps@gmail.com editor@ijlbps.org

PROFILE OF ANTIBIOTIC USE AT THE HEALTH CENTRE

¹Dr. B. Deepak,²Mr. Ch. Rajendra Prasad,³Mrs. J. Shireesha

^{1,2,3}Assistant Professor

¹Department of Pharmacy Practice

^{2,3}Department of Pharmacology

Vaagdevi Institute of Pharmaceutical Sciences, Bollikunta, Warangal. Telangana.

ABSTRACT

Context: Excessive use of antibiotics is a prevalent issue in healthcare, resulting in needless drug expenditure, increased susceptibility to side effects, and the emergence of antibiotic resistance. Unsafe and ineffective therapy, illness progression, and increased health care expenses are all consequences of inappropriate prescribing practices. This study's objective was to evaluate the University of Gondar referral hospital's outpatient pharmacy department's antibiotic prescribing practices using World Health Organization prescribing indicators in Gondar, Northwest Ethiopia. Methods: A cross-sectional study that was retrospective was carried out. Prescription and prescription registration books kept at the pharmacy store were used to gather prescription data for a year. Prescription indicators from the World Health Organization and the International Network of Rational Use of Drugs were used to gauge rational drug use, with a particular emphasis on antibiotic prescribing trends. Version 20 of SPSS was used to examine the data that was gathered. Findings and Discussion: Out of 600 patient interactions, 968 medications were administered. There were 1.6 drugs on average each contact. The proportion of consultations where injections and antibiotics were administered was 6.3% and 69.7%, respectively. The most often given antibiotics were amoxicillin (28.5%),

ciprofloxacin (12%), and metronidazole (11.1%). 95.3% of prescribed medications came from the list of necessary pharmaceuticals, and 96% came from generic names. The World Health Organization's recommended standard for the rate of antibiotic prescriptions was not followed, but there were minor variations from the standard in the areas of polypharmacy, injectable prescribing patterns, brand name usage, and prescriptions for medications from the National Essential Drugs List. In order to prevent the inappropriate use of antibiotics and prevent additional consequences, interventions targeted at changing the prescribing patterns for antibiotics must be put into place.

Keywords: antimicrobial resistance, prescribing indicators, prescribing patterns, and antibiotics

1. INTRODUCTION

Antibiotics are medications that can destroy or inhibit the growth of bacteria by either selectively killing or inhibiting the development of disease-causing bacteria.^{1,2} Currently, they are the most commonly prescribed drugs in hospitals worldwide. Antibiotics play a pivotal role in combating disease and maintaining health especially in developing countries where infectious diseases are still a big challenge. However, in recent years their benefit is facing a great challenge due to the

emergence of antibiotic resistance. Currently, it is found that many microbes have become resistant to the most commonly available and effective first line agents mainly due to inappropriate prescribing practices.³⁻⁵ The consumption of antibiotics has increased worldwide with most of this occurring in low- and middle-income countries.⁶ Antibiotics prescription varies significantly between countries as shown by 1 out of 2 hospitalized patients receiving antibiotics in Africa and Asia while 1 out of 3 patients in Europe receives antibiotics.⁷ In many African countries, including Ethiopia, it is a common practice to prescribe multiple medications in a single prescription paper, which is referred to as polypharmacy, and in the current study it refers to using 5 and more medications in a single prescription paper. A study conducted on 26 primary care facilities in Cameroon showed that from the total 30 096 prescription papers reviewed, 11035 (36.7%) had at least 1 antibiotic prescription.³ Another study conducted in Lesotho showed that the average number of medicines prescribed per prescription was 3.8 of which antibiotics constitute 37.6%.⁸ Many studies conducted in Ethiopia also showed that antibiotics were among the most commonly prescribed groups of medications. Antibiotic resistance is a global health crisis and is one of the greatest challenges for public health and affects both developing and developed countries.¹²⁻¹⁴ Inappropriate prescribing habits lead to ineffective and unsafe treatment, worsening/prolongation of diseases and cost increments due to further complication of diseases and the need for further treatment. Inappropriate prescribing also reduces the quality of medical care and leads to wastage of resources.

In developing countries antibiotics are prescribed for 44% to 97% of hospitalized

patients often unnecessarily or inappropriately.^{16,17} It is estimated that half of all medicines in Africa are used inappropriately including two third of antibiotics.^{18,19}

Many studies conducted in Ethiopia have shown that most of the bacteria that cause infections to human beings and animals have developed considerable degree of resistance to commonly used first-line antibiotics.^{7,21} The current study was aimed at assessing antibiotics prescribing pattern at university of Gondar comprehensive and specialized hospital using the WHO prescribing indicators.

2. METHODS AND MATERIALS

Study Area and Period

The study was conducted at the University of Gondar comprehensive and specialized hospital outpatient Pharmacy Department. The hospital is a governmental hospital located in Gondar town, central Gondar zone, Amhara regional state, Ethiopia. Gondar is located at 727 km away from Addis Ababa; capital city of the country and 185 km from Bahirdar, the capital city of the region. University of Gondar comprehensive and specialized hospital is one of the oldest and respected teaching hospitals in the country currently serving more than 4.1 million people of the central Gondar zone and people from neighboring regions. The study was conducted from March 5, 2019 to June 10, 2019 state, Ethiopia. Gondar is located at 727 km away from Addis Ababa; capital city of the country and 185 km from Bahirdar, the capital city of the region. University of Gondar comprehensive and specialized hospital is one of the oldest and respected teaching hospitals in the country currently serving more than 4.1 million people of the central Gondar zone and people from

neighboring regions. The study was conducted from March 5, 2019 to June 10, 2019.

Study Design

A retrospective cross-sectional study was conducted. Data were collected using a prequalified check list. World Health Organization/International Network of Rational Use of Drugs (WHO/INRUD) prescribing indicators with their standard values were utilized to measure rational use of drugs with due focus on antibiotics prescribing patterns.

Population

Source Population. All prescription papers (23000) that were dispensed between March 1, 2018 and March 30, 2019 at the Outpatient Pharmacy Department of University of Gondar Comprehensive and Specialized Hospital.

Sample Size Determination and Sampling Procedure

WHO's "How to investigate drug use in health facilities" recommends at least 600 encounters to be included in a cross-sectional survey involving a drug use survey.²² Because of financial constraints to include more encounters, 600 prescription papers for a 1-year period were studied per the WHO recommendation.²² Specific prescription papers were selected using systematic random sampling method, using prescription numbers as a reference for a sampling frame. The total number of prescription papers dispensed between March 1, 2018 and March 30, 2019 were 23000 and from thus dividing this with the number of the sample size "600" gave us the sampling interval "38." Every prescription paper has its unique identification number and thus all of the prescriptions dispensed during the study

period were arranged in ascending order and every "38th" prescription paper was selected.

Data Processing and Analysis

The collected data were entered in to SPSS version 20 and analyzed. As the study was conducted in a hospital setting, all the drugs dispensed require a valid prescription paper. In the statistical analysis, frequencies, averages/means, and percentages were calculated. Percentage of encounters in which an antibiotic was prescribed was calculated to measure the overall use of commonly overused and costly forms of drug therapy. It was calculated by dividing the number of patient encounters in which an antibiotic was prescribed by the total number of encounters surveyed (600), multiplied by 100.

Data Quality Assurance

Pretest of the data collection tool was conducted on 30 prescription papers in order to assess the adequacy of the questionnaire to determine the parameters in question. The collected data were checked for completeness, accuracy, and consistency at the end of each day.

Definition of Terms

Antibiotics include penicillins, other antibacterials, antileprosy drugs, anti-infective dermatological drugs, anti-infective ophthalmological agents, anti-diarrhoeal drugs with streptomycin, neomycin, nifluroxazide, or combinations.²² Combination of drugs indicates 2 or more drugs that are prescribed for a given health condition. For example, triple therapy for *Helicobacter pylori*-induced peptic ulcer is counted as one.

3. Results Completeness of the Prescription

Demographic data of patients (age and sex) as well as date of prescription were mentioned in all the prescriptions reviewed. However, none of the prescriptions contained patient weight and height data. Drug-related information, such as name of the drug, strength, frequency and duration of treatment, prescriber, and dispenser information were

Table 1. Sociodemographic Characteristics of Sampled Patients at University of Gondar Comprehensive and Specialized Hospital Outpatient Pharmacy Department Between March 2018 and March 2019.

Characteristics	Frequency	Percentage
Gender		
Male	242	40
Female	358	60
Age, years		
<5	46	7.7
5-14	60	10
15-30	302	50.3
31-44	87	14.5
45-64	85	14.2
>65	19	3.2

Table 2. Summary Report of Number of Drugs per Encounter at University of Gondar Comprehensive and Specialized Hospital Outpatient Pharmacy Department Between March 2018 and March 2019.

Number of Drugs per Encounter	Frequency	Average (%)
One drug	335	55.8
Two drugs	168	28.0
Three drugs	91	15.3
Four drugs	6	1.0

also completely mentioned in all of the prescriptions reviewed. The diagnosis of ailments was recorded in 212 (35.3%) of the prescription papers.

Drug-Related Outcomes

As shown in Table 1, the majority of the prescriptions (358, 60%), were prescribed

for females. Majority of the patients were between the ages of 15 and 30 years. A total of 968 medicines were prescribed with an average number of drugs per encounter found to be 1.6. Out of all prescriptions, 335(55.8%) of them had only 1 drug per prescription while 6 prescriptions contained 4 drugs (Table 2). The total number of encounters prescribed with antibiotics and injection were 418(69.6%) and 38(6.3%) respectively. About 95% of the drugs (923, 95.3%), were prescribed from the essential drug list of Ethiopia.²³ The details are shown in Table 3.

Table 3. Summary Report of Prescribing Indicators at University of Gondar Comprehensive and Specialized Hospital Outpatient Pharmacy Department Between March 2018 and March 2019.

Prescribing Indicators	Total Number of Drugs	Average (%)	WHO Standard Value
1. Average number of drugs per encounter	968	1.6	(1.4-1.8)
2. Percentage of encounters with antibiotics	418	69.6	(20-34.8)
3. Percentage of encounters with injection	38	6.3	(13.4-24.1)
4. Percentage of drugs prescribed by generic name	929	96	100
5. Percentage of drugs from essential drug list	923	95.3	100




Figure 1. Total number of antibiotics per prescription at University of Gondar Comprehensive and Specialized Hospital Outpatient Pharmacy Department between March 2018 and March 2019.

By antibiotics category, penicillin was the most frequently prescribed (217, 38.2%) category of antibiotics followed by macrolides (84, 15%; Figure 2).

By specific types of antibiotics, amoxicillin was the most frequently prescribed antibiotic (162, 28.6%), followed by ciprofloxacin (66, 12%) and metronidazole (63, 11.1%; Table 4).

Among the total 212 prescriptions for which the diagnosis was written, 162 contained antibiotics. Among the 162 prescriptions, gastrointestinal tract infections (44, 28%) were the most common diagnosis for frequently prescribed antibiotics followed by lower respiratory tract infections (30, 19.4%; Table 5).

Majority of antibiotics were prescribed by oral route (476, 84%) followed by parenteral route (39, 4%). Regarding the dosage form of antibiotics, capsule (255, 45%) and tablets (231, 40.7%) were the most commonly prescribed dosage forms.

Discussion Completeness of Prescriptions

All the 600 prescriptions contain patient information (name, age, and sex). This was greater than a study conducted in Addis Ababa and earlier in Gondar in which only 25% and 85% of the prescriptions contain patient information (age and sex), respectively.^{9,6} However, only 35.3% of the prescriptions contained patient diagnosis information, which is much lower than the standard value (100%). But this finding was better than a similar study conducted earlier in Gondar in 2011, which was only 1.4%.⁶ Regarding the drug-related information in this study, all the prescriptions contain the correct name, strength, duration, frequency, and dosage form of the drugs (100%). This was better than a study conducted earlier in Gondar, which was 79%.⁶ This could be attributed to improvements in prescribing and dispensing practice by health professionals, which could in




Figure 2. Most commonly prescribed classes of antibiotics at University of Gondar Comprehensive and Specialized Hospital Outpatient Pharmacy Department between March 2018 and March 2019.

Table 4. Summary of Most Commonly Prescribed Antibiotics at University of Gondar Comprehensive and Specialized Hospital Outpatient Pharmacy Department Between March 2018 and March 2019.

Common Antibiotics	Frequency	Percentage
Amoxicillin	162	28.6
Ciprofloxacin	67	12
Metronidazole	63	11
Azithromycin	59	10.4
Ceftriaxone	37	6.5
Cephalexin	36	6.3
Amoxicillin + clavulanate	33	5.8
Doxycycline	32	5.6
Cloxacillin	26	4.5
Clarithromycin	25	4.4
Trimethoprim-Sulfamethoxazole	16	2.8
Norfloxacin	13	2.2

turn be due to continuous professional developments and revolutionary practices executed by the hospital and specially the pharmacy department.

Average Number of Drugs per Encounter

In the current study, the average number of drugs per encounter was 1.6, which is within the range recommended Table 5. Summary of Common Diagnosis for Frequently Prescribed Antibiotics at University of Gondar Comprehensive and Specialized Hospital Outpatient Pharmacy Department Between March 2018 and March 2019.

Diagnosis	Frequency	Percentage
1. Gastrointestinal infection	44	27.5
2. Lower respiratory infection	30	19.0
3. Sexually transmitted diseases	29	18.1
4. Upper respiratory infection	27	17.0
5. Urinary tract infection	21	13
6. Skin and soft tissue infection	10	6.1

by WHO (1.6-1.8).²⁴ This finding was lower than a similar study conducted in Addis Ababa, Hawassa, Ghana, and earlier in Gondar, in which the average number of drugs per encounter was 2.0, 1.9, 3.5, and 1.77, respectively.^{6,9,11,25} The lower the number of drugs prescribed per encounter; it is a positive sign of good prescribing practice. It reduces polypharmacy and in turn minimizes disease complication due to drug-drug interactions and adverse drug reaction.

In this study the most common indications for antibiotic prescription was respiratory tract infections (57, 36%) followed by gastrointestinal tract infections (44, 27.5%) and sexually transmitted diseases (29, 18.125%). This result was consistent with a similar study conducted in Addis Ababa in which respiratory tract infections were the most common indications for antibiotics (24.5%).⁹ This could be due to the fact that respiratory tract infections such as tonsillitis and pharyngitis, which usually occur due to problems in personal hygiene, are very common in Ethiopia and antibiotics are the mainstream treatment, which makes them being frequently prescribed.

Use of Antibiotics

In the current study, percentage of antibiotics per prescription was found to be 69.6%, which is much higher than the ideal value recommended by WHO (20-26.8).²² This finding was much higher than earlier study conducted in Gondar on drug use practice in which only 29.3% of the

prescriptions contained an antibiotic.⁶ It was also higher than similar studies conducted in Addis Ababa and southern Ethiopia, in which antibiotics constitute 38% and 58.1%, respectively

In this study, penicillins were the most frequently prescribed classes of antibiotics (38.5%) followed by macrolides (15%) and fluoroquinolones (14%). This finding was consistent with a similar study conducted in Addis Ababa and Dessie, in which penicillins (51.9%) were the most commonly prescribed groups of antibiotics followed by fluoroquinolones (18.3%).^{9,10} A similar finding was reported by a study in Kenya in which penicillins (46.9%) followed by cephalosporins (45.8%) were the most commonly prescribed classes of antibiotics.²⁷ A study conducted in Bangladesh, however, reported that cephalosporins were the most frequently prescribed groups of antibiotics followed by macrolides.²⁸ By specific antibiotics, amoxicillin (28.5%) followed by ciprofloxacin (12%) and metronidazole (11.1%) were the most frequently prescribed antibiotics in the current study. This finding was consistent with similar study conducted in Ghana, in which amoxicillin (22.5%) followed by ciprofloxacin (18.4%) being the most frequently prescribed antibiotics.²⁵ Amoxicillin was also found to be among the commonly prescribed antibiotics in studies conducted in Addis Ababa and Dessie.^{9,10} But this was different from a study done in Kenya, in which ceftriaxone was the most frequently prescribed antibiotics (39.7%) followed by benzyl penicillin (29%).²⁷ This might be due to regional variation in bacterial susceptibility/resistance, prescribing habit and the difference in prevalence of infectious diseases in different countries.

4. CONCLUSION

The current study's antibiotic prescribing pattern deviates from the WHO's suggested guideline. This means that, in order to prevent the unfavorable effects of administering antibiotics inappropriately, ongoing interventional techniques and routine audits are required at all levels of healthcare.

However, despite a little divergence from the WHO's suggested standard values, polypharmacy, injectable prescribing, brand name usage, and prescribing medications on the Essential Drugs List were not shown to be significant issues in this study.

Recommendations

The hospitals need to encourage systematic monitoring of antibiotic use through institution of programs to promote rational drug use. The hospital's Drug and Therapeutic Committee should be strengthened to control antibiotic use and regularly provide clinicians with up-to-date information regarding rational antibiotic prescribing practice. The ministry of health should update and disseminate national standard treatment guidelines, encourage targeted, problem-based in-service educational programs and regular continuing education for health professionals.

REFERENCES

1. European Centre for Disease Prevention and Control. Factsheet for experts—antimicrobial resistance. <https://www.ecdc.europa.eu/en/antimicrobial-resistance/facts/factsheets/experts>. Published 2008. Accessed October 7, 2019.
2. Leekha S, Terrell CL, Edson RS. General principles of antimicrobial therapy. *Mayo Clin Proc.* 2011;86:156-167. doi:10.4065/mcp.2010.0639
3. Chem ED, Anong DN, Akoachere JKT. Prescribing patterns and associated factors of antibiotic prescription in primary health care facilities of Kumbo East and Kumbo West Health Districts, North West Cameroon. *PLoS One.* 2018;13:e0193353. doi:10.1371/journal.pone.0193353
4. Kimang'a AN. A situational analysis of antimicrobial drug resistance in Africa: are we losing the battle? *Ethiop J Health Sci.* 2012;22:135-143.
5. Llor C, Bjerrum L. Antimicrobial resistance: risk associated with antibiotic overuse and initiatives to reduce the problem. *Ther Adv Drug Saf.* 2014;5:229-241. doi:10.1177/2042 098614554919
6. Admassie E, Begashaw B, Tesfaye WH. Assessment of drug use practices and completeness of prescriptions in Gondar university teaching referral hospital. *Int J Pharm Sci Res.* 2013;4:265-275.
7. Drug Administration and Control Authority of Ethiopia; Management Sciences for Health, Strengthening Pharmaceutical Systems. Antimicrobials use, resistance and containment baseline survey syntheses of findings. <http://apps.who.int/medicinedocs/en/m/abstract/Js18057en/>. Published August 2009. Accessed October 7, 2019.
8. Ntšekhe M, Hoohlo-Khotle N, Tlali M, Tjipura D. Antibiotic prescribing pattern at six hospitals in Lesotho. <http://apps.who.int/medicinedocs/documents/s21028en/s21028en.pdf>. Published July 2011. Accessed October 7, 2019.
9. Worku F, Tewahido D. Retrospective assessment of antibiotics prescribing at public primary healthcare facilities in Addis

Ababa, Ethiopia. *Interdiscip Perspect Infect Dis.* 2018; 2018:4323769. doi:10.1155/2018/4323769

10. Getachew E, Aragaw S, Adissie W, Agalu A. Antibiotic prescribing pattern in a referral hospital in Ethiopia. *Afr J Pharm Pharmacol.* 2013;7:2657-2661.

11. Desalegn AA. Assessment of drug use pattern using WHO prescribing indicators at Hawassa University Teaching and Referral Hospital, south Ethiopia: a cross-sectional study. *BMC Health Serv Res.* 2013;13:170. doi:10.1186/1472-6963-13-170

12. Goossens H, Ferech M, Stichele RV, Elseviers M; ESAC Project Group. Outpatient antibiotic use in Europe and association with resistance: a cross-national database study. *Lancet.* 2005; 365:579-587. doi:10.1016/s0140-6736(05)17907-0

13. Center for One Health Research, University of Washington. Antimicrobial resistance. <https://deohs.washington.edu/cohr/antimicrobial-resistance-0>. Published 2019. Accessed October 7, 2019.

14. World Health Organization. Antibiotic resistance: multicountry public awareness survey. <https://www.who.int/drugresistance/documents/baselinesurvey-nov2015/en/>. Published November 2015. Accessed October 7, 2019.

15. World Health Organization. Promoting rational use of medicines: core components—WHO Policy Perspectives on Medicines, No. 005, September 2002. <http://apps.who.int/medicinedocs/en/d/Jh3011e/>. Published September 2002. Accessed October 7, 2019.