



ISSN 2395-650X

**International Journal** of  
Life Sciences Biotechnology Pharma Sciences

IJLBPS



[www.ijlbps.org](http://www.ijlbps.org)

E-mail: [editorijlbps@gmail.com](mailto:editorijlbps@gmail.com) [editor@ijlbps.org](mailto:editor@ijlbps.org)

# Investigation of the Phytochemical Components, Antimicrobial Actions, and Impact of *Pycanthus Angolensis* Weld Warb Exudate on Corneal Ulcers in Rabbits (Myristicaceae)

S.Umarani <sup>1</sup>, Hanumanthu <sup>2</sup>,Kandarpalle Rahul <sup>3</sup>,

## Abstract

*For the purpose of treating corneal ulcers, a combination of 1 part Anchomonas difformis, 1 part Cyrtosperma senegalense, and 2 parts Pycanthus angolensis has been proposed. In accordance with claims made in traditional medicine, this research will assess the phytochemical components, antibacterial activity, and efficacy of a combination of these exudates in treating corneal ulcers. The method included collecting fresh exudates in separate containers. After they were dried off, the phytochemical components of each residue were examined. We further investigated the antibacterial properties of P. angolensis exudates by studying their impact on rabbits with chemically produced ocular ulcers. The results showed that the exudates of A. angolensis and C. senegalensis contained only reducing sugars. Tannins, flavonoids, and reducing sugars were the bioactive components found in P. angolensis exudates. Additionally, this demonstrated antibacterial action against the tested species. Within ten days of therapy, it cured the corneal ulcers in rabbits that had been caused by NaOH. In conclusion, the corneal ulcers in rabbits were cured by the exudates of P.angolensis, which included bioactive components and showed antibacterial activity. Even without A. angolensis and C. senegalensis exudates, its traditional usage for healing corneal ulcers seems reasonable.*

*Important terms: rabbits, Pycanthus angolensis, ocular ulcers, exudates.*

## INTRODUCTION

*Pycanthus angolenses* Welw Warb (Myristicaceae) is also known as “African nutmeg or false nutmeg”<sup>1</sup> The Nigerian Local names of the plant are, Abakang (Ibibio) Akwa mili (Ibo) Abora (Itsekiri), Akamo, Akujaadi (Yoruba), Nupe (Kpokgi), and Abaororo (Urhobo). Others are: Etena (Cameroon), and Loioka (Zaire)<sup>1</sup> It is a forest tree of about 30.5 m high 2.45 m in girth. The bark is grey, longitudinally fissured, flaking in patches in old trees and exudes reddish coloured juice. It is widely used for ethnomedical purposes<sup>2</sup> An infusion of the bark is reported to be effective in the treatment of leprosy<sup>3</sup> and for

purification of breast milk in Guinea. A terpenoid quinine with potential use in treatment of Type 2 diabetes was isolated from the plant<sup>4</sup> Herbal medicine practioners in some parts of Delta State, Nigeria, claim that a 1: 1: 2 mixture of exudates of *Anchomonas difformis*: *Cyrtosperma senegalense*: and *Pycanthus angolensis* respectively is used for treatment of corneal ulcers. (Personal communication with Tega Ikuegbvweha, the second-named author). This work was done to verify this claim.

*Assistant professor <sup>1,2,3</sup>,  
Department of Pharmacy,  
Samskruti College of Pharmacy,  
Kondapur (V), Ghatkesar (M) Medchal Dist, Telangana, India.*

## EXPERIMENTAL

### *Plants exudates.*

Parts of *Anchomonas difformis*, and *Pycanthus angolensis* plants were collected from Ikpoba Hill, Benin City and *Cryptosperna senegalense* was from Urhobo by the second named-author. They were identified by Alhaji A. Abubakar a Taxonomist formally with the Department of Pharmacognosy, Faculty of Pharmacy, University of Benin. Exudates from each plant was collected into a separate bottle, reduced to dryness on the boiling water bath and the residue subjected to the tests.

### *Phytochemical tests:*

Phytochemical tests for bioactive constituents were carried out on portions of the residual material using standard phytochemical procedures<sup>5- 8</sup>.

### *Fehling's test for reducing sugars (in glycosides):*

The residue was re-dissolved in water on the water bath. To 2ml of the solution, in the test tube was added, 1ml each of Fehling's solutions A and B. The mixture was shaken and heated in a water bath for 10min. The colour obtained was recorded. A brick-red precipitate indicates reducing sugar.

### *Frothing test for saponins:*

Water extract was obtained by boiling on the water bath . The extract was transferred into a test tube and shaken vigorously then was left to stand for 10 mins and the result noted. A thick persistent froth indicates saponins.

### *Borntrager's test for anthraquinone derivatives:*

Chloroform extract of the material was obtained by boiling on the water bath. To 2 mls of this extract, 1ml of dilute (10 %) ammonia was added and the mixture was shaken. Any colour change was recorded. A pink-red colour in the ammoniacal (lower) layer shows anthracene derivatives. *Test for Flavonoids* Water extract of the sample was reduced to dryness on the boiling water bath. The residue was treated with dil. NaOH, followed by addition of dilute HCl, solubility and colour were noted. A yellow solution with NaOH, which turns colourless with dil HCl confirms flavonoids.

### *Ferric chloride solution test for tannins:*

Water extract was treated with 15 % ferric chloride test solution. The resultant colour was noted. A blue colour indicates condensed tannins, a green colour indicated hydrolysable tannins.

### *Colour tests for alkaloids:*

500 mg of plant material was extracted with 500 mls of methanol for 20 minutes, on a water bath The extract was then filtered off and allowed to cool. This extract was dispensed in 2 ml of portions into four different test tubes. Either the Dragendorff's or Hager's or Mayer's or Wagner's alkaloidal reagent was added to each tube and the presence or absence of colours of any precipitates were noted in each test tube. *Keller- Kiliiani test* (for de-oxy sugars in cardiac glycosides): Methanol extract was obtained and the extract reduced to dryness. 50 mg of this was dissolved in 2 ml chloroform. H<sub>2</sub>SO<sub>4</sub> was added to form a layer and the colour at interphase recorded. Brown ring at interphase is characteristic of deoxysugars in cardenolides. The results showed the presence of only reducing sugars in *A. difformis* and *C. senegalense* but showed that *P. angolensis* contained reducing sugars, as well as tannins and flavonoid glycosides. These are constituents that could have biological activity, hence the biological tests were carried out on the *P. angolensis* exudates only.

**Table 1:** Phytochemical constituents of exudates of *Pycanthus angolensis*

Test	Observation	Inference
Test for reducing sugars in glycosides (Fehling's solutions test)	Blue colour of Fehling's solutions turned brick red ppt.	Presence of reducing sugar
Frothing test for Saponins	Only bubbles seen. No frothing	Absence of Saponins
Borntrager's reagent test for anthracene derivatives	Same shade of pink as the exudates	Absence of Anthracene derivatives
Keller Kiliiani test for deoxysugars in cardiac glycosides	No brown ring colour	Absence of deoxysugars, no cardiac glycosides.
FeCl <sub>3</sub> solution test for Tannins	Blue-black ppt	Hydrolysable Tannins present
Test for flavonoids NaOH Test	Flavonoids dissolved giving yellow colour. solution turns colorless with HCl	Flavonoids present.
HCl test Tests with Dragendorff's, Hager's, Mayer's, Wagner's alkaloidal Reagents for Alkaloids	No Coloured ppt. with any reagent	Absence of Alkaloids

**Table 2:** Antibacterial activities of exudates of *P. angolensis* and reference drugs on test bacterial organisms

Treatment (ug/disc)	Organisms and diameters of Zones of inhibition (mm)			
	<i>S. aureus</i>	<i>Strep. pyrogens</i>	<i>E. coli</i>	<i>P. aeruginosa</i>
Exudates: 50	8.2	10.4	10.5	6.1
100	12.6	16.6	14.3	7.5
250	19.4	21.5	15.2	10.6
Ampicillin 20	25.8	28.8	18.4	12.2
Gentamicin 20	18.3	23.7	27.9	14.1

**Table 3:** Effects of exudates of *P. angolensis* on chemical-induced corneal ulcers in rabbits


Parameter	Ocular state of ulcerated and treated rabbits	
	1 <sup>st</sup> Day	10 <sup>th</sup> Day
Status of eyelid	Badly inflamed	No Inflammation
Lacrimation	Very severe	Normal
Discharge	Nil	Nil
Corneal Tissue	Badly ulcerated	Healed



A  
Normal left eye rabbit



B  
Right eye on 1<sup>st</sup> day of  
ulcer induction



Right eye of B on 10<sup>th</sup> day  
of treatment with exudates

**Fig 1:** Result of Treatment of induced corneal ulcers with *P. angolensis* exudates

**Test Organisms:**

Standard Laboratory strains of the organisms were obtained from the University of Benin Teaching Hospital and sub cultured in the Laboratory of Pharmaceutical Microbiology, University of Benin. The organisms used were *Escherichia coli* (Gram-negative), *Pseudomonas aeruginosa* (Gram-negative) *Staphylococcus aureus* (Gram-positive) and *Streptococcus pyrogens* (Gram-positive). The bacterial cultures were maintained on nutrient agar (Oxoid Ltd). Prior to use, they were sub-cultured into nutrient broth and incubated overnight at 37 OC.

**Antibacterial tests**

The activities of exudates on the organisms were evaluated using the paper disc diffusion method (9). 100 mg of dried residue was redissolved in 10 ml normal saline and from this stock solution volumes

equivalent to 50, 100 and 250 µg/disc were spotted on antibiotic sterile assay paper discs (Whatman No. 1 (0.6cm diameter). A blank containing normal saline and another containing 20 µg/disc of a particular reference drug (Ampicillin or gentamicin) were also set up. The discs were allowed to air-dry and then placed on pre-sterilized agar plates (10 cm diameter) containing 10 ml of Agar Muller Hinton 3.5% medium, and spread with 10 µl of a standardized inoculum ( $5 \times 10^5$  cells/ml) of the test bacterium. Plates were incubated for 24 hr. at 37°C and zones of inhibition measured (mm). Experiments were set up in four replicates. The results are presented in Table 2.

### ***Effect of exudates on corneal ulcers in rabbits.***

#### ***Animals:***

Rabbits, 300-450 g body weight were bought from Aduwawa in Benin City. They were maintained on potato leaves, elephant grass and mature unripe pawpaw fruits and water for a period of 8 weeks for them to acclimatize.

#### ***Animal Care:***

They were de-wormed by 0.1ml ivermectin SuperR subcutaneously (sc) administered. Also to improve their immunity and check infection and stress, keproceryl and vitalite were given orally. ***Induction of ulcer.*** Animals were anaesthetized with ketamin injection (sc) 2 drops of 4% Novesea tropical anesthetic agent instilled into the cornea of right eyes of 2 rabbits, (A) followed 20 mins later with 2 drops of 1M solution of NaOH instilled into the eyes. The 2 left eyes (B) were left as control. The 2 animals showed ulcerated cornea (A) and so were used for the test. The dried residue was re-dissolved in normal saline to make a stock of 2% w/v from this, 2 drops were daily instilled into the right ulcerated eyes (A). The extract was administered 3 times daily for nine consecutive days following induction. Animals were allowed free food and water. During this period, observations were made for any signs of toxicity, infection and discharge, status of the ulcerated cornea, any healing. Photograph of the ulcerated, non-ulcerated and treated eyes were taken on the 1st day of treatment and on the 10th day of experimentation. The photographs are enclosed. (Fig1)

### **RESULTS AND DISCUSSION**

Exudates of *A. difformis* and *C. senegalense*, contained only reducing sugars. Exudate of *P. angolensis* contained reducing sugars, flavonoids and tannins. In the antimicrobial test, (Table 2) 50 µg/ml

exudate had least activity against all the organisms. It was most active against *Strep pyrogens*. Though 250 µg/ml showed significant activity on the organisms, these were less than those of the reference drugs. The induced ulcers were healed within nine days (Fig 1). There was no sign of toxicity in the animals that received the exudates. Phenolic compounds are generally noted for their antimicrobial activities.<sup>5,8</sup> Specifically, antimicrobial activities of tannins and flavonoids of *Allium cepa* <sup>10</sup> and *Thymus serpyllum*<sup>11</sup> have been reported. These phenolic constituents present in the exudates could then be responsible for its antibacterial effect as noted in this work. In addition, tannins act externally as water proof to the external layers of the exposed tissue since they precipitate proteins, thus protecting the underlying layers. They also have vasoconstriction effect on small superficial or exposed vessels. By limiting fluid losses and by preventing external aggressions, tannins enhance tissue regeneration in case of superficial wounds or burns<sup>8,10</sup>. This further strengthens the healing activity of the exudates. Thus the healing effect of exudates could also be attributed to the above actions of Flavonoids and tannins.

### **CONCLUSION**

This work has identified the phytochemical constituents in exudates of *A. difformis*, *C. senegalense* and *P. angolensis*. The antibacterial activities of exudates of *P. angolensis*, though less than the activities of the reference drugs, ampicillin and gentamicin, has been demonstrated. The *P. angolensis* exudates alone was active, it healed corneal ulcers within nine days of treatment, in contrast to the claim that the mixture of exudates of the three plant materials exerts the effect.

### **REFERENCES**

1. Bep O. Medicinal Plants in Nigeria. Published as a Private Edn The Nigerian College of Arts, Science and Technology Ibadan. 1960: 3-20.
2. Akendengue B and Louis AM. Medicinal Plants used by the Masanga People in Gabon. *J. Ethnopharmacol* 1994: 41 193-200.
3. Vasileva, B. Medicinal Plants of Guinea. Conakry Republique de Guinea Masco USSA. Moscow University 1969: p 1.
4. Jeamecheung LJ, Yevish EM, Clarke JP, Tsai J, Priscilla Lopresca, J Ubillas RP, Fort DM, Hector RF, King SR, Mendez CD, Gerald DJ, Hector RF, King SR, Mendez CD, Jolad SD, Rewoven GM. Novel type quinones isolated from *Pycnanthus angolensis* of potential utility in the treatment of Type 2 diabetes. *J. Pharmacology and Therapeutics*, 1999; 2: 529-534.

5. Bruneton J. Pharmacognosy, Phytochemiistry and Medicinal Plants. Intercept Ltd Londers, New York. 2nd Edn. p. 386.
6. Sofowora A. Medicinal Plants and Traditional Medicine in Africa. John Wiley and Sons Ltd. Chichester. 1982, pp 142-145.
7. Harborne JB. Phytochemical Method: A guide to Modern Techniques of Plants Analysis. 2<sup>nd</sup> Edn. Chapman and Hall New York. 1983.
8. Evans WC. Trease and Evans Pharmacognosy 13th Edn E.LBS with Bailliere Tindall. 1989 pp. 388, 480, 502, 535, 546.
9. Bauer AW, Kirby WMM, Sherris JC, Turck M. Antibiotic susceptibilty testing by a standardized single disk method. *Am J.Clin Pathol.* 1966; 45: 493-496.
10. Izo AA, Cario GD, Briscards D. Biological screening of Italian medicinal plants for antibacterial activity. *Phytother. Res.* 1995. 9: 281-286
11. Janssen AM, Scheffer JJC, Baerheim SA. Antimicrobial activity of essential oils. A 1996- 1998 Literature Review. *Plants Med.* 1987. 53. 395-398.