



ISSN 2395-650X

**International Journal** of  
Life Sciences Biotechnology Pharma Sciences

IJLBPS



[www.ijlbps.org](http://www.ijlbps.org)

E-mail: [editorijlbps@gmail.com](mailto:editorijlbps@gmail.com) [editor@ijlbps.org](mailto:editor@ijlbps.org)

## The effectiveness of modifying Park's approach in treating strabismus in youngsters

G.Thanusha <sup>1</sup>, G.Suresh <sup>2</sup>, L.Sunil <sup>3</sup>, Bishwajit Patowary <sup>4</sup>,

### Abstract

**Goal:** Examine if modified Park's approach works well for strabismus patients' kids.

**Methods:** From January 2019 to December 2021, a total of 120 patients were recruited at the Anhui Provincial Children's Hospital in Anhui, China, in the Department of Ophthalmology. Each of the two groups—the study and the control—contained sixty patients. The research group underwent a modified Park's approach, which featured an intermuscular membrane incision and a conjunctiva two-layer suture method, whereas the control group got rectus muscle adjustment suture using a normal incision. Numerous factors were evaluated, including patient satisfaction, tear film performance, and perioperative indications.

**Findings:** There was a substantial reduction in intraoperative blood loss, surgical time, and hospital stay for the study group ( $p < 0.01$ ). Additionally, it had a substantially lower corneal staining score ( $p < 0.01$ ), a significantly greater Schirmer's time, and a tear film break-up time (TFBUT) time. The study group's satisfaction level was much greater than the control group's ( $p < 0.05$ ). Clinical effectiveness was also higher (91.67%) than in the control group (83.33%). Furthermore, compared to the control group (11;  $p < 0.05$ ), the study group showed a considerably decreased incidence of complications (5).

**In conclusion,** enhanced Park's approach results in excellent effectiveness and a decreased incidence of problems while also improving perioperative indicators, tear film function, and satisfaction level. This implies that it might be a good alternative to the standard care given to kids with exotropia. However, in order to prove that this treatment approach is better, long-term follow-up data will be needed.

**Keywords:** conjunctiva, two-layer suture, intramuscular membrane, Park's method, and strabismus

### INTRODUCTION

Strabismus, characterized by misaligned extra ocular muscles, can result in a deviation in eye position. It is a relatively common condition among adolescents, with a prevalence of around 4 %. Treatment of strabismus is time-consuming and often leads to psychological stress for both patients and their families [1,2]. Surgical correction is currently the main approach, but it carries the risk of complications such as corneal exposure and surface damage, which may affect tear film function and impact surgical outcomes and patient satisfaction [3,4]. As a result, safeguarding the cornea during surgery and reducing postoperative complications have become key priorities in clinical

practice. In recent years, rectus recession has emerged as a preferred surgical option for strabismus treatment. It offers advantages such as shorter operation time, improved visual field during surgery, and fewer postoperative side effects [5]. However, this technique is not without issues, including eyelid scarring and conjunctival wounds [6]. In comparison, modified Park's technique has gained popularity in strabismus surgery due to its smaller conjunctival incisions, reduced postoperative discomfort, and minimal aesthetic impact [7]. In

**Assistant professor** <sup>1,2,3,4</sup>,  
**Department of Pharmacy,**  
**Samskruti College of Pharmacy,**  
**Kondapur (V), Ghatkesar (M) Medchal Dist, Telangana, India.**

traditional Chinese medicine (TCM), strabismus in children is classified as ophthalmoplegia or visual divergence, often attributed to blocked meridians caused by *wind*, *phlegm*, and *stasis*. This obstructs the flow of *qi* and blood while depriving tendons and muscles of nourishment and relaxation, resulting in oblique eyes. Intermittent external strabismus is mostly associated with *qi* deficiency, resulting from lack of positive *qi*, weakened external guard consolidation, and deficiency in veins and ligaments. Hence, this study incorporates the use of *Buzhong Yiqi* decoction as a treatment approach. This study investigated the efficacy of modified Park's technique in the treatment of strabismus in children.

## METHODS

### General patient data

This research was conducted at Anhui Provincial Children's Hospital in Hefei, Anhui Province, China, comprising 120 children diagnosed with strabismus. Patients were randomly assigned using an online web-based tool (<http://www.randomizer.org/>) to two groups namely; study and control groups. An independent research assistant, not involved in participant screening or evaluation, managed the randomization process. The original calculation for sample size determined that 60 patients in each group would be sufficient to detect a 3-point difference between two groups in a two-sided significance test, with a statistical power of 0.8 and an alpha error level of 0.05.

All patients and their guardians received comprehensive information and provided informed consent by signing respective documentation before participating in the study which adhered to the ethical standards outlined in the Declaration of Helsinki [8]. The study received approval from the Ethics Committee of Anhui Provincial Children's Hospital, China (approval no. CHA18239474).

### Inclusion criterion

Patients who met diagnostic criteria for strabismus as specified in the 9th edition of ophthalmology [9].

### Exclusion criteria

Patients with abnormal eye movements or A-V signs, nystagmus, vertical strabismus, a history of eye surgery or trauma, or any other ocular diseases, surgical intolerance, abnormal immune function, refractive interstitial opacity, or neurological diseases were excluded from this study.

### Interventions

Patients were in the supine position and received general anesthesia. In preparation for surgery, the

conjunctival sac was rinsed with 0.3 % povidone-iodine. Compound epinephrine was applied to the conjunctival surface to constrict capillaries and minimize intraoperative blood loss.

For medial rectus surgery, an arcuate incision was made 5 mm away from the cornea in the nasal region. For lateral rectus surgery, an arcuate incision was made 7 mm away from the cornea in the temporal region. The bulbar conjunctiva was cut open, and the rectus muscle was secured in place using a strabismus hook. Thereafter, bulbar conjunctiva and fascia were dissected to fully expose the rectus muscle. A double-loop suture was performed using a 6 - 0 absorbable suture, approximately 2 mm from the endpoint of rectus muscle. The medial or lateral rectus muscle was then cut about 1 mm posterior to its end to allow for recession. Suture was advanced obliquely at the posterior edge of the insertion point of rectus muscle, emerging from the anterior edge. Needle was passed through the superficial sclera at the end of the rectus muscle. Configuration of the suture resembled a "V", with its tip facing the cornea, and the amount of muscle recession was measured. The suture was tightened without ligating according to the preoperative suspension muscle design. The conjunctiva was secured with a silk thread. Following operation, tobramycin and dexamethasone ophthalmic ointment were applied to the conjunctival sac. If necessary, the suture was adjusted 1-2 days after surgery based on eye position. In control group, rectus muscle adjustment suture was performed using the conventional incision method. A 5 – 10 mm bulbar conjunctival fornix incision was made, and the medial or lateral rectus muscle was secured using a strabismus hook. Intermuscular membrane and ligamentum temperatum were dissected. After performing a double-loop suture using 6 - 0 absorbable sutures, extraocular muscles were shortened according to preoperative design, and conjunctival incision was sutured. Both groups of patients received oral dexamethasone tablets (5 mg/day) for three consecutive days. Levofloxacin and prapofen eye drops were administered four times daily for two consecutive weeks.

Additionally, both groups received treatment with *Buzhong Yiqi* decoction, which included 12 g each of *Codonopsis radix* and *Astragali radix*, 9 g each of *Atractylodis macrocephalae* rhizoma, *Rehmanniae radix*, *Angelicae sinensis* radix, *Bupleuri radix*, *Scrophulariae radix*, and *Cimicifugae* rhizoma, and 6g each of tangerine peel, *Ophiopogonis radix*, and

liquorice root. The herbs were decocted with water and administered every other day for six months.

### Evaluation of parameters/indices

#### Postoperative indicators

Amount of blood loss during surgery, duration of surgical procedure, and length of hospital stay was recorded.

#### Efficacy

Efficacy was classified as cured (following treatment, symptoms vanished, and eye position was restored, with degree of strabismus decreasing to less than  $5\Delta$ ), effective (a noticeable improvement in symptoms, and degree of strabismus dropped to between  $5\Delta$  and  $10\Delta$  after treatment), and ineffective (despite treatment, there was no significant improvement in symptoms) [10].

#### Tear film function

Schirmer's test procedure involved placing a sterile test paper below the patient's eye in the conjunctival sac. Patient was instructed to look straight ahead for a few seconds and then keep their eyes closed. After 5 mins, the paper strip was collected, and the length of moistened area was measured. In the tear film break-up time (TFBUT) test, sodium fluorescein was applied to the conjunctival surface, and eye was examined using a slit lamp. Lamp was switched to a cobalt blue filter, and patient was instructed to blink once and keep their eyes open. A dark spot indicating a dry area appeared after the patient blinked. Tear film break-up time (TFBUT) was calculated as the time interval between the last blink and appearance of the first dark spot. For corneal fluorescein staining score, a solution of sodium fluorescein was placed on patient's cornea, and eye was observed under cobalt blue light to determine staining pattern. Cornea was divided into quadrants, and each quadrant was assigned a score from 0 to 3. Total score across all quadrants ranges from 0 to 12 with lower scores indicating better tear film function.

#### Satisfaction with surgery

After a 28-day post-surgery period, a survey was administered to evaluate patient satisfaction. The survey included questions regarding the efficacy of surgery and any discomfort experienced. Each question was assigned a score of 1 to 5. A score of 80 – 100 indicated a very high level of satisfaction, 60 – 80 indicated satisfaction while < 60 indicated dissatisfaction.

#### Complications

Occurrences of complications, which included conjunctivitis, corneal edema, diplopia and scar adhesion were recorded.

#### Statistical analysis

Data processing was conducted using Statistical Packages for Social Sciences (SPSS 23.0).

Enumeration data were presented as N (%) and analyzed using Chi-square test. Normally distributed measurement data were represented as mean  $\pm$  standard deviation (SD). Comparison of means between two groups was carried out using chi-square F-test. Data showing chi-square differences were tested using independent samples t-test, while data with non-chi-square differences were analyzed using independent samples t-test. Intra-group pre- and post-comparisons were performed using paired samples t-test.  $P < 0.05$  was considered statistically significant.

## RESULTS

### Patient data

There was no statistical difference in clinical data between study and control groups ( $p > 0.05$ ; Table 1).

**Table 1:** Comparison of general data (mean  $\pm$  SD, N = 60)

Characteristic	Study group	Control group	t/ $\chi^2$	P-value
Sex (M/F)	43/37	42/38	1.667	0.197
Age (years)	6.75 $\pm$ 1.22	6.34 $\pm$ 1.45	0.049	0.961
Duration of disease (months)	2.42 $\pm$ 0.79	2.14 $\pm$ 1.17	2.064	0.356
Left eye	36	41		
Right eye	44	39	0.054	0.816
Degree of strabismus	46.01 $\pm$ 10.22	45.82 $\pm$ 9.65	0.123	0.902

### Perioperative indicators

Perioperative indices in study group showed significant improvement compared to control group. ( $p < 0.05$ ; Table 2).

#### Tear film function

Following intervention, the use of modified Park's technique led to significantly improved tear film function, when compared to rectus muscle adjustment suture.

Data was significantly reduced in both groups after intervention, and tear film function was significantly better in study group compared to control group ( $p < 0.05$ ; Table 3).

#### Satisfaction level

Satisfaction level in study group was significantly higher compared to control group ( $p < 0.05$ ; Table 4).

#### Clinical efficacy

Patients in study group exhibited higher treatment efficacy rate compared to control group ( $p < 0.05$ ) (Table 5).

#### Incidence of complications

Study group had lower incidence rate of complications compared to control group ( $p < 0.05$ ) (Table 6).

**Table 2:** Perioperative indices (mean  $\pm$  SD, N = 60 in each group)

Group	Intraoperative blood loss (mL)	Duration of surgery (min)	Length of hospital stay (days)
Study	8.11±2.12	22.14±5.02	5.87±1.53
Control	13.21±3.07	35.23±6.41	7.86±2.35
χ <sup>2</sup> /t	9.326	10.632	5.476
P-value	<0.01	<0.01	<0.01

**Table 3:** Comparison of tear film function (mean ± SD, N = 60 in each group)

Group	Schirmer's test (mm.5min)		TFBUT (s)		Corneal staining score (points)	
	Before intervention	After intervention	Before intervention	After intervention	Before intervention	After intervention
Study	5.62±1.21	10.42±1.54*	5.58±1.23	8.12±1.39*	8.27±1.32	4.25±1.27*
Control	3.67±1.05	11.33±1.35*	3.64±1.07	9.13±1.41*	8.22±1.31	6.16±1.24*
T	0.042	7.322	0.042	7.632	0.058	7.322
P-value	0.974	<0.01	0.954	<0.01	0.942	<0.01

\*P < 0.05 vs. before intervention

**Table 4:** Comparison of satisfaction (N = 60 in each group)

Group	Very satisfied	Satisfied	Dissatisfied	Overall satisfaction
Study	32	17	1	59
Control	14	26	10	50

**Table 5:** Efficacy of treatment (N = 60 in each group)

Group	Cure	Effective	Ineffective	Total efficacy (%)
Study	26	29	5	91.67
Control	18	32	10	83.33

**Table 6:** Incidence of complications (N= 60 in each group)

Group	Conjunctivitis	Corneal edema	Diplopia	Scar adhesions	Total
Study	1	2	1	1	5
Control	4	3	2	2	11
χ <sup>2</sup>					4.234
P-value					0.035

## DISCUSSION

Strabismus is commonly associated with craniocerebral injury and neurological diseases

[11,12]. It typically manifests before age 5, causing alignment issues and vision loss as children struggle to align their eyes correctly [13]. Vision training, along with appropriate eye surgery, aids in the recovery of eyesight, with corrective surgery being the primary management method for strabismus in current practice [13-15].

For children with external strabismus, the root cause often lies in an internal deficiency of *middle-qi* and lack of glory of *Ying* and blood. This leads to deficiency of *yin* and blood and excessive *yang* activity, resulting in intermittent and sudden eye strabismus. Main clinical symptoms include *qi* deficiency, with strabismus occurring after physical exhaustion or loss of mental focus. *Buzhong Yiqi* decoction, containing *Astragalus* radix benefits *qi*, *Codonopsis* radix and *Atractylodes macrocephalae* rhizoma strengthen and nourishes spleen and *middle-qi*, tangerine Peel regulates *qi*, *Angelicae sinensis* radix tonifies blood, *Cimicifugae* rhizoma and *Bupleuri* radix elevate and strengthens *middle-qi*, *Ophiopogonis* radix nourishes *Yin* and blood, are used as a combination treatment option.

The modified technique of Park's incision, combined with two layers of intermuscular and conjunctival sutures, contributes to the restoration of tertiary visual function in both eyes of children with external strabismus. It also promotes the attainment of functional eye position after surgery. Compared to control group, study group showed lower perioperative scores, which is attributed to accurate lesion localization and smaller conjunctival incision using modified Park's technique. These factors significantly reduced intraoperative bleeding, operative time, and postoperative complications. The less invasive nature accelerated healing, shortened hospital stays reduced pain and treatment costs. Additionally, study group exhibited improved tear film function when compared to control group. Thus, modified Park's technique was less damaging to aqueous and mucus layers of tear film. Furthermore, this procedure minimizes conjunctival scarring, nerve damage and irritation in postoperative period, leading to enhanced tear film stability and recovery [16-18].

Study group surpassed control group in terms of treatment efficacy, complication rate and satisfaction with surgery primarily due to smaller conjunctival incision and reduced postoperative pain. Park's technique, originally developed by Dr Marshall Parks, an American ophthalmologist in 1968, eliminates the need for tissue dissection anterior to muscle insertion, thereby reducing nerve damage. This may be considered as a therapeutic option in future.

### Limitations of this study

This current study has some limitations. Firstly, the sample size used was relatively small, which may limit the generalizability of findings. Secondly, lack of long-term follow-up data did not establish durability of treatment outcomes over an extended period.

## CONCLUSION

Modified Park's technique is highly efficacious for the treatment of strabismus in children. This technique minimizes surgical wounds and reduces the length of surgical procedures and hospital stays. As a result, patients undergoing this technique experience a lower incidence of complications and a higher level of satisfaction. Thus, modified Park's technique may be considered a preferred surgical option for treating strabismus in children following collection of long-term follow-up data to establish durability of treatment in the future.

## REFERENCES

- Heydarian S, Abbasabadi MM, Khabazkhoob M, Hoseini-Yazdi H, Gharib M. Vision abnormalities in children and young adults with cerebral palsy; a systematic review. *Semin Ophthalmol* 2022; 37(4): 471-479.
- Kulp MT, Ciner E, Ying GS, Candy TR, Moore BD, Orel-Bixler D. Vision screening, vision disorders, and impacts of hyperopia in young children: outcomes of the vision in preschoolers (VIP) and vision in preschoolers - hyperopia in preschoolers (VIP-HIP) studies. *Asia-Pacific J Ophthalmol* 2022; 11(1): 52-58.
- Su H, Fu J, Wu X, Sun A, Zhao BW, Hong J. Comparison of Botulinum toxin type A with surgery for the treatment of intermittent exotropia in children. *BMC Ophthalmol* 2022; 22(53): 1-8
- Temeltürk RD, Koçer AM, Yasar HH. Psychosocial improvements in children and their parents following strabismus surgery. *J Pediatr Ophthalmol Strabismus* 2022; 59(4): 261-268.
- Nwanaji-Enwerem JC, Gateman T, Whitecross S, Whitman MC. First visit characteristics associated with future surgery in intermittent exotropia. *J Binocul Vis Ocul Motil* 2022; 72(1): 22-28.
- Shen T, Kang Y, Lin XM, Wu HP, Yan JH. Newly developed abnormal head position and secondary esotropia after strabismus surgery for children with intermittent exotropia. *Int Ophthalmol* 2022; 42(10): 2965-2972.
- Zhang XY, Fu K, Jiang YZ. Effects of limbal trapezoidal conjunctival flap incision and modified Parks incision on tear film function and postoperative complications in patients with strabismus. *Int J Ophthalmol* 2021; 21(04): 618 -622.
- World Medical Association. World Medical Association Declaration of Helsinki: Ethical principles for medical research involving human subjects. *JAMA* 2013; 310: 2191-2194.
- Yang PZ, Fan XQ. *Ophthalmology*. 9th edition. Beijing: People's Health Publishing House, 2018: 247-253.
- Tian L, Shen ML, Gao DJ. Clinical observation of Park's incision continuous suture without knots in children with concomitant strabismus under the microscope. *China J Tradit Chin Med* 2021; 31(02): 103-106.